



RISK AND POLICY ANALYSIS UNIT

Working Paper
RPA 18-02
April 24, 2018

Quasi Maximum Likelihood Analysis of High Dimensional Constrained Factor Models

Kunpeng Li

Qi Li

Lina Lu

Quasi Maximum Likelihood Analysis of High Dimensional Constrained Factor Models*

Kunpeng Li[†], Qi Li[‡] and Lina Lu^{§¶}

April 24, 2018

Abstract

Factor models have been widely used in practice. However, an undesirable feature of a high dimensional factor model is that the model has too many parameters. An effective way to address this issue, proposed in a seminal work by Tsai and Tsay (2010), is to decompose the loadings matrix by a high-dimensional known matrix multiplying with a low-dimensional unknown matrix, which Tsai and Tsay (2010) name the constrained factor models. This paper investigates the estimation and inferential theory of constrained factor models under large- N and large- T setup, where N denotes the number of cross sectional units and T the time periods. We propose using the quasi maximum likelihood method to estimate the model and investigate the asymptotic properties of the quasi maximum likelihood estimators, including consistency, rates of convergence and limiting distributions. A new statistic is proposed for testing the null hypothesis of constrained factor models against the alternative of standard factor models. Partially constrained factor models are also investigated. Monte carlo simulations confirm our theoretical results and show that the quasi maximum likelihood estimators and the proposed new statistic perform well in finite samples. We also consider the extension to an approximate constrained factor model where the idiosyncratic errors are allowed to be weakly dependent processes.

Key Words: Constrained factor models, Maximum likelihood estimation,
High dimensionality, Inferential theory.

JEL #: C13, C38.

*We would like to thank the guest editors and two anonymous referees for their insightful comments that greatly improve our paper. We also thank Jushan Bai for the helpful comments on an early version of this paper. Kunpeng Li gratefully acknowledges the financial support from China NSF No.71722011 and No.71571122. Qi Li's research is partially supported by China NSF project No.71133001 and No 71601130.

[†]International School of Economics and Management, Capital University of Economics and Business, Fengtai District, Beijing, 100070, China. Email: kunpenglithu@126.com.

[‡]Corresponding author: Qi Li, Email: qi-li@tamu.edu. Department of Economics, Texas A&M University, College Station, TX 77843, USA,

[§]Federal Reserve Bank of Boston, the Risk and Policy Analysis Unit, Department of Supervision, Regulation, and Credit, 600 Atlantic Avenue, Boston, MA 02210, USA. Email: lina.lu@bos.frb.org

[¶]The views expressed in the paper are those of the authors and should not be interpreted as reflecting the views of the Federal Reserve Bank of Boston, or any other person associated with the Federal Reserve System.

1 Introduction

With the rapid development of data collection, storing and processing techniques in computer science, econometricians and statisticians now face large dimensional data setups more often than ever before. A challenge along with the appearances of large data is how to extract useful information from data, or put differently, how to effectively conduct dimension reduction on data. Factor models are proved to be an effective way to perform this task. Over the last three decades, the literature has witnessed wide applications of factor models in many economics disciplines. In finance, Conner and Korajczyk (1986, 1988) and Fan, Liao and Shi (2014) use factor models to measure the risk and performance of large portfolios. In macroeconomics, Geweke (1977) and Sargent and Sims (1977) use dynamic factor models to identify the source of primitive shocks. In labor economics, Heckman, Stixrud and Urzua (2006) use factor models to capture unobservable personal abilities. In international economics, Kose, Otrok and Whiteman (2003) use multilevel factor models to separate global business circles, regional business circles and country-specific business circles. Large dimensional factor models are also used in a variety of ways to deal with strong correlations, see e.g., Fan, Liao and Mincheva (2011) and Fan, Liao and Mincheva (2013), among others.

A standard factor model can be written as

$$z_t = Lf_t + e_t, \quad t = 1, 2, \dots, T,$$

where $z_t = (z_{1t}, \dots, z_{Nt})'$ is a vector of N variables at time t , L is an $N \times r$ loadings matrix, f_t is an r -dimensional vector of factors and e_t is an N -dimensional vector of idiosyncratic errors. The traditional (classical) factor analysis assumes that N is fixed and T is large. This assumption runs counter to usual shape of large dimensional data sets, in which N is usually comparable to or even greater than T (Stock and Watson (2002)). Recent literature contributes a lot to the asymptotic theory with N comparable to or even greater than T . Bai and Ng (2002) propose several information criterions to determine the number of factors in a large- N and large- T environment. Under a similar setup to Bai and Ng (2002), Stock and Watson (2002) prove that the principal components (PC) estimates are consistent in approximate factor models of Chamberlain and Rothschild (1983). Bai (2003) moves forwards along the work of Stock and Watson (2002) and gives the asymptotic representations of the PC estimates of loadings, factors and common components. Doz, Giannone and Reichlin (2012) consider the maximum likelihood (ML) method and prove the average consistency of the maximum likelihood estimates (MLE). Bai and Li (2012, 2016) use five different identification strategies to eliminate the rotational indeterminacy from asymptotics and give limiting distributions of the MLE. Fan, Liao and Wang (2014) propose a new projected principal component method to more accurately estimate the unobserved latent factors.

A potential problem in high dimensional factor models is that too many parameters

are estimated within the model, which makes it difficult to analyze and interpret the economic implications of the estimates. However, if the space of the loading matrix is spanned by a low dimension matrix, this problem can be much ameliorated. In this paper, following Tsai and Tsay (2010), we address this problem by considering the following constrained factor model

$$z_t = M\Lambda f_t + e_t,$$

where M is a *known* $N \times k$ matrix with rank k and Λ is a $k \times r$ *unknown* loadings matrix with rank r . We assume $r < k \leq C$ for some generic constant C . In the above specification, M consists of the bases of the loading matrix. The underlying true loadings are a weighted average of these bases associated with the weights matrix Λ , which are the parameters of interests. The number of loading parameters now is kr instead of Nr . So the number of parameters is greatly reduced.

Our work is closely related to Tsai and Tsay (2010) who were the first to consider constrained factor models. This paper differs from Tsai and Tsay (2010) in several dimensions. First, although Tsai and Tsay propose using PC and ML methods to estimate constrained factor models, their asymptotic analysis focuses only on the PC method. They obtain convergence rates of the PC estimates. As a comparison, we investigate asymptotics of the ML method and derive the convergence rates and limiting distributions of the MLE. Given the limiting distributions, one can easily construct $(1 - \alpha)$ -confidence intervals if prediction is the target of interest, or use t -test or F -test to conduct statistical inferences on the underlying parameter values if hypothesis testing is the purpose. Second, Tsai and Tsay consider the setup that k is large (but still smaller than N). In this paper, we instead assume that k is fixed^①. In our viewpoints, assuming a fixed k is of practical and theoretical interests. In some typical examples, the parameter k is interpreted as the number of groups or categories, according to which the variables are classified (see Tsai and Tsay (2010)). This value is usually not large in real data. Therefore, a fixed- k assumption is adopted in this paper. Furthermore, in constrained factor models, a large k leads to a larger number of parameters being estimated. The estimation accuracy is reversely linked with k for a given sample size. When k is large, the benefit of constrained factor models against standard factor models becomes weak, which makes constrained factor less attractive in practice. Third, an importantly related issue in constrained factor models is on conducting valid model specification check on the presence of matrix M . Tsai and Tsay consider the traditional likelihood ratio test to perform this task. But the traditional likelihood ratio test is designed under fixed- N and large- T setup, which conflicts to large- N and large- T scenarios. In this paper, we propose new statistics for testing model specifications that are applicable to the large- N and large- T setups.

^①Our analysis can be extended to the case of a large k . But for this case, deriving the limiting distribution of the MLE is very challenging since the matrix Λ is high-dimensional.

The rest of the paper is organized as follows. Section 2 provides more empirical examples of the constrained factor model. Section 3 introduces the model and lists the assumptions needed for the subsequent analysis. Section 4 delivers the consistency and limiting distribution results of the MLE. Section 5 considers testing issues within constrained factor models. Section 6 considers a partially constrained factor model and presents the asymptotic properties of the MLE for this model. Section 7 presents the Expectation-Maximization (EM) algorithm to implement the QML estimation. Section 8 conducts Monte Carlo simulations to investigate the finite sample performance of the MLE and to study the empirical size and power of the proposed model specification test. In Section 9, we relax Assumption B to allow for the idiosyncratic errors to have a more general weakly dependence structure. Section 10 concludes the paper. All technical contents are delegated to several appendices.

2 Motivating Applications

The well-known equilibrium arbitrary pricing theory (APT) implies that the observed assets returns can be expressed into a linear factor structure, see Ross (1976), Conner and Korajczyk (1988) among others. This motivates the use of the following factor model

$$r_{it} = \sum_{j=1}^r l_{ij} f_{jt} + e_{it}$$

to study the performance of portfolios, where r_{it} is the excess return of the i th security at time t , f_{jt} denotes the j th risk premium at time t and l_{ij} the beta coefficient of the j th risk premium for security i . However, as pointed out by Rosenberg (1974), the common movements among the assets returns may be related with the individual characteristics. Such characteristics include capitalization and book-to-price ratios as suggested in Fama and French (1993), momentum as in Carhart (1997), own-volatility as in Goyal and Santa-Clara (2003). Let x_{ip} denote the observed p th characteristic of the i th security. Rosenberg (1974) considers the specification

$$l_{ij} = \sum_{p=1}^k x_{ip} \lambda_{pj} + v_{ij}, \quad \text{or} \quad L = M\Lambda + V,$$

where $M = (x_{ip})_{N \times k}$ is the observed characteristics matrix. Rosenberg's specification is very close to the one studied in this paper. With a slight modification, the analysis in this paper can easily be extended to cover the Rosenberg's model.

A limitation of Rosenberg's specification is that the factor betas are assumed to be linear functions of the observed characteristics, which is overly restrictive in practice. To accommodate this concern, Conner and Linton (2007) and Conner, Hagmann and Linton (2012) consider the following nonparametric specification

$$l_{ij} = g_j(x_{ij}).$$

where $g_j(\cdot)$ is an unknown smooth function. Conner, Haggmann and Linton (2012) apply their model to a real dataset and indeed find that the factor betas are nonlinear functions of the characteristics. However, an undesirable feature in these two papers is that the estimation of the model involves an iterative procedure between the factors and unknown functions, which is formidable to many applied researchers. To address this issue, we instead consider using a series of polynomial functions to approximate the unknown function $g_j(\cdot)$. More specifically, we consider approximating the function $g_j(\cdot)$ by all the polynomial functions with power less than q , i.e.,

$$g_j(x) \approx \lambda_{j0} + \lambda_{j1}x + \cdots + \lambda_{jq}x^q. \quad (2.1)$$

Given this, the model now can be written as $L = M\Lambda$ with

$$M = \begin{bmatrix} 1 & x_{11} & x_{11}^2 & \cdots & x_{11}^q & \cdots & \cdots & x_{1r} & x_{1r}^2 & \cdots & x_{1r}^q \\ 1 & x_{21} & x_{21}^2 & \cdots & x_{21}^q & \cdots & \cdots & x_{2r} & x_{2r}^2 & \cdots & x_{2r}^q \\ \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_{N1} & x_{N1}^2 & \cdots & x_{N1}^q & \cdots & \cdots & x_{Nr} & x_{Nr}^2 & \cdots & x_{Nr}^q \end{bmatrix}$$

and

$$\Lambda = \begin{bmatrix} \lambda_{10} & \lambda_{11} & \cdots & \lambda_{1q} & 0 & \cdots & 0 & \cdots & \cdots & 0 & \cdots & 0 \\ \lambda_{20} & 0 & \cdots & 0 & \lambda_{21} & \cdots & \lambda_{2q} & \cdots & \cdots & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \ddots & \ddots & \vdots & \ddots & \vdots \\ \lambda_{r0} & 0 & \cdots & 0 & 0 & \cdots & 0 & \cdots & \cdots & \lambda_{r1} & \cdots & \lambda_{rq} \end{bmatrix}'.$$

The above model can be viewed as a special case of the constrained factor model with some zero restrictions imposed on Λ . The model considered here maintains the nonlinear function feature of Conner and Linton (2007) and Conner, Haggmann and Linton (2012) but the computational burden has been much reduced. A primary issue related with our method is whether the approximation (2.1) is good enough. This work can be partially addressed by the W statistic proposed in Section 5.

Constrained factor models have other applications. Tsai and Tsay (2010) apply constrained factor models to analyze stock returns where the stocks can be classified into different sectors. They specify the constraint matrix M consisting of orthogonal and binary vectors. In another application, they implement constrained factor models to study the interest-rate yield curve, where the columns of the matrix M are specified to denote the level, slope and curvature feature of interest rates. Matteson et al. (2011) use constrained factor models to forecast the hourly emergency medical service call arrival rates by specifying the constraints on the factor loadings based on the prior information of the pattern of the call arrivals. Similar approach is adopted in Zhou and Matteson (2015) to model the ambulance demand by incorporating covariate information as constraints on the factor loadings.

3 Constrained Factor Models

Let N denote the number of variables and T the sample size in the time dimension. We consider the following constrained factor model

$$z_t = M\Lambda f_t + e_t, \quad (3.1)$$

where $z_t = (z_{1t}, z_{2t}, \dots, z_{Nt})'$ is an N -dimensional vector of explanatory variables at time t ; M is a specified $N \times k$ (known) matrix with rank k ; Λ is the $k \times r$ loading matrix of rank r ; $f_t = (f_{1t}, f_{2t}, \dots, f_{rt})'$ is a vector of r latent common factors; e_t is an N -dimensional vector of idiosyncratic disturbances and is independent of f_t . Throughout the paper, we assume $k \geq r$. If $k < r$, we can simply consider the linear regression $z_t = Mf_t^* + e_t$ with $f_t^* = \Lambda f_t$. The model effectively becomes a factor model with k (when $k < r$) factors.

Our analysis is based on similar assumptions used in standard factor models, see Bai and Li (2012) for the asymptotic analysis of the MLE for standard high dimensional factor models. The symbol C appearing in the following assumptions denotes a generic constant. Our assumptions include:

Assumption A: $\{f_t\}$ is a sequence of fixed constants with $\bar{f} = \sum_{t=1}^T f_t = 0$. Let $M_{ff} = \frac{1}{T} \sum_{t=1}^T f_t f_t'$ be the sample variance of f_t . There exists an $\bar{M}_{ff} > 0$ (positive definite) such that $\bar{M}_{ff} = \lim_{T \rightarrow \infty} M_{ff}$.

Assumption B: The idiosyncratic error term e_{it} is independent across the i index and the t index with $E(e_t) = 0$, $E(e_t e_t') = \Sigma_{ee} = \text{diag}(\sigma_1^2, \sigma_2^2, \dots, \sigma_N^2)$ and $E(e_{it}^8) \leq C$ for all i and t , where $e_t = (e_{1t}, e_{2t}, \dots, e_{Nt})'$ is the N -dimensional vector of idiosyncratic errors at time t .

Assumption C: The underlying values of parameters satisfy that

- C.1 $\|\Lambda\| \leq C$ and $\|m_j\| \leq C$ for all j , where m_j is the transpose of the j th row of M .
- C.2 $C^{-2} \leq \sigma_j^2 \leq C^2$ for all j , where $\sigma_j^2 = E(e_{jt}^2)$ is defined in Assumption B.
- C.3 Let $P = \Lambda' M' \Sigma_{ee}^{-1} M \Lambda / N$, $R = M' \Sigma_{ee}^{-1} M / N$. We assume that $P_\infty = \lim_{N \rightarrow \infty} P$ and $R_\infty = \lim_{N \rightarrow \infty} R$ exist. In addition, $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \sigma_i^{-4} (m_i \otimes m_i)(m_i' \otimes m_i') = V_\infty$ exists. Here P_∞ , R_∞ and V_∞ are some positive definite matrices.

Assumption D: The estimator of σ_j^2 for $j = 1, \dots, N$ takes value in a compact set: $[C^{-2}, C^2]$. Furthermore, M_{ff} is restricted to be in a set consisting of all semi-positive definite matrices with all elements bounded in the interval $[-C, C]$, where C is a large positive constant.

Assumption A requires that factors are sequences of fixed constants. The random factors can be dealt with in a similar way under some suitable moment conditions. Assumption B is commonly imposed in classical factor models. It can be relaxed to allow for cross-sectional and temporal heteroskedasticities and correlations, see Bai and Li (2016)

for a related development in this direction. Assumption C requires that underlying values of parameters are in a compact set, which is standard in econometric literature. Assumption D requires that some parameter estimates take values in a compact set. This assumption is often made when dealing with highly nonlinear objective function, see Jennrich (1969). Our objective function is highly nonlinear.

Similar to the case of a standard factor model, a constrained factor model has an identification problem. To see this, for any invertible $r \times r$ matrix B , we have

$$\Lambda f_t = \Lambda B \cdot B^{-1} f_t = \Lambda^* f_t^*.$$

with $\Lambda^* = \Lambda B$ and $f_t^* = B^{-1} f_t$. To separate (Λ, f_t) from (Λ^*, f_t^*) , we impose the following identification condition.

Identification condition (abbreviated by IC hereafter):

IC1 $\Lambda' (\frac{1}{N} M' \Sigma_{ee}^{-1} M) \Lambda = P$, where P is a diagonal matrix whose diagonal elements are distinct and arranged in a descending order.

IC2 $M_{ff} = \frac{1}{T} \sum_{t=1}^T f_t f_t' = I_r$.

Our identification strategy is similar to IC3 in Bai and Li (2012). It is known that this identification strategy identifies the loadings and factors up to a column sign, see Bai and Li (2012) for a detailed discussion on this issue. To eliminate such a problem in our theoretical analysis, we follow Bai and Li (2012) to treat as part of the identification condition that the estimators and the underlying values of loadings matrix have the same column signs. In practice, the sign problem causes no troubles in empirical analysis.

We use the following discrepancy function between M_{zz} and Σ_{zz} as our objective function

$$\mathcal{L}(\theta) = -\frac{1}{2N} \ln |\Sigma_{zz}| - \frac{1}{2N} \text{tr}[M_{zz} \Sigma_{zz}^{-1}], \quad (3.2)$$

where $\theta = (\Lambda, \Sigma_{ee})$, $M_{zz} = T^{-1} \sum_{t=1}^T z_t z_t'$ and $\Sigma_{zz} = M \Lambda \Lambda' M' + \Sigma_{ee}$. This discrepancy function has the same form as a likelihood function when f_t are independently and normally distributed with mean zero and variance I_r , see Bai and Li (2012) for details. In the current paper, the factors are assumed to be fixed constants in Assumption A, the above discrepancy function is therefore not a likelihood function. Nevertheless, we still call the maximizer of the above function as a quasi MLE or MLE for simplicity. Specifically, the MLE $\hat{\theta} = (\hat{\Lambda}, \hat{\Sigma}_{ee})$ is defined as

$$\hat{\theta} = \underset{\theta \in \Theta}{\text{argmax}} \mathcal{L}(\theta),$$

where Θ is the parameters space such that any interior point of it satisfies Assumption D and the identification condition IC. The input parameters include Λ and Σ_{ee} . In a constrained factor model, we only need to estimate kr loadings instead of Nr loadings (the number of parameters in a standard factor model). Therefore, the number of parameters

is greatly reduced. Taking derivatives with respect to Λ and Σ_{ee} , we obtain the following first order conditions:

$$\hat{\Lambda}' M' \hat{\Sigma}_{zz}^{-1} (M_{zz} - \hat{\Sigma}_{zz}) \hat{\Sigma}_{zz}^{-1} M = 0; \quad (3.3)$$

$$\text{diag}(\hat{\Sigma}_{zz}^{-1}) = \text{diag}(\hat{\Sigma}_{zz}^{-1} M_{zz} \hat{\Sigma}_{zz}^{-1}), \quad (3.4)$$

where $\hat{\Lambda}$ and $\hat{\Sigma}_{ee}$ denote MLE of Λ and Σ_{ee} , respectively, and $\hat{\Sigma}_{zz} = M \hat{\Lambda} \hat{\Lambda}' M' + \hat{\Sigma}_{ee}$. We note that the above two first order conditions are only used in deriving the asymptotic properties of the MLE. One does not need to solve the above nonlinear equations to obtain the MLE. Instead, we can implement the EM algorithm to compute the MLE. Details are given in Section 7.

4 Asymptotic properties of the MLE

In this section, we investigate the asymptotic properties of the MLE. The following proposition shows that the MLE is consistent.

Proposition 4.1 (Consistency) *Let $\hat{\theta} = (\hat{\Lambda}, \hat{\Sigma}_{ee})$ be the MLE that maximizes (3.2). Then under Assumptions A-D, together with IC, when $N, T \rightarrow \infty$, we have*

$$\hat{\Lambda} - \Lambda \xrightarrow{p} 0; \quad \frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2 \xrightarrow{p} 0.$$

In high dimensional factor analysis, the loadings and variances of idiosyncratic errors are high-dimensional. The consistencies have to be defined under some chosen norms, see Stock and Watson (2002), Bai (2003), Doz, Giannone and Reichlin (2012) and Bai and Li (2012, 2016). In constrained factor models, due to the presence of matrix M , the loading matrix Λ is low-dimensional. So its consistency is defined in the elementwise sense. But for the variances of idiosyncratic errors, they are still high-dimensional. Their consistency is therefore defined by $\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2$, which can be written as $\frac{1}{N} \|\hat{\Sigma}_{ee} - \Sigma_{ee}\|^2$. So the chosen norm is the Frobenius norm adjusted with the matrix dimension.

Given the consistency results, we have the following theorem on convergence rates of the MLE.

Theorem 4.1 (Convergence rates) *Under the assumptions of Proposition 4.1, we have*

$$\hat{\Lambda} - \Lambda = O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T}\right), \quad \frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2 = O_p\left(\frac{1}{T}\right).$$

According to Theorem 4.1, the convergence rate of $\hat{\Lambda}$ is $\min(\sqrt{NT}, T)$, which is faster than the \sqrt{T} -convergence rate of estimated loadings in standard factor models. This result is plausible since in a constrained factor model, we use NT observations to estimate kr loadings. This is in contrast with a standard factor model, where we use NT observations to estimate Nr loadings.

To present the asymptotic representation of the MLE, we introduce some notation.

Let

$$\mathbb{D}_1 = \begin{bmatrix} 2D_r^+ \\ \mathcal{D}[(P \otimes I_r) + (I_r \otimes P)K_r] \end{bmatrix}, \quad \mathbb{D}_2 = \begin{bmatrix} 2D_r^+ \\ 0_{\frac{1}{2}r(r-1) \times r^2} \end{bmatrix}, \quad \mathbb{D}_3 = \begin{bmatrix} 0_{\frac{1}{2}r(r+1) \times r^2} \\ \mathcal{D} \end{bmatrix},$$

and

$$\mathbb{B}_1 = K_{kr}[(P^{-1}\Lambda') \otimes \Lambda] + R^{-1} \otimes I_r - K_{kr}(I_r \otimes \Lambda)\mathbb{D}_1^{-1}\mathbb{D}_2[(P^{-1}\Lambda') \otimes I_r],$$

$$\mathbb{B}_2 = K_{kr}(I_r \otimes \Lambda)\mathbb{D}^{-1}\mathbb{D}_3(\Lambda \otimes \Lambda)', \quad \Delta = \mathbb{B}_2 \frac{1}{N} \sum_{i=1}^N \frac{1}{\sigma_i^6} (m_i \otimes m_i)(\kappa_{i,4} - \sigma_i^4),$$

where $P = \frac{1}{N}\Lambda'M'\Sigma_{ee}^{-1}M\Lambda$, $R = \frac{1}{N}M'\Sigma_{ee}^{-1}M$, $\kappa_{i,4} = E(e_{it}^4)$, m_i is the transpose of the i th row of matrix M , K_{uv} is the commutation matrix such that for any $u \times v$ matrix B , $K_{uv}\text{vec}(B) = \text{vec}(B')$; and K_r is defined to be K_{rr} . $D_r^+ = (D_r'D_r)^{-1}D_r'$ is the Moore-Penrose inverse matrix of the r -dimensional duplication matrix D_r , \mathcal{D} is the matrix such that $\text{veck}(B) = \mathcal{D}\text{vec}(B)$ for any $r \times r$ matrix B , where $\text{veck}(B)$ is the operation which stacks the elements below the diagonal of the matrix B into a vector. Given matrix P , we can easily calculate the matrix \mathbb{D}_1 and its inverse. For example, let $P = \text{diag}(1, 2, 3)$ ($r = 3$ in this case), then

$$\mathbb{D}_1 = \begin{bmatrix} 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2 \\ 0 & 1 & 0 & 2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2 & 0 & 3 & 0 \end{bmatrix}, \quad \mathbb{D}_1^{-1} = \begin{bmatrix} 0.5 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 1.5 & 0 & 0 & 0 & 0 & -0.5 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0.5 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 3 & 0 & 0 & 0 & -1 \\ 0 & 0 & -0.5 & 0 & 0 & 0 & 0 & 0.5 & 0 \\ 0 & 0 & 0 & 0 & -2 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0.5 & 0 & 0 & 0 \end{bmatrix}.$$

Now we state the asymptotic result of $\hat{\Lambda}$.

Theorem 4.2 (Asymptotic representation) *Under assumptions of Theorem 4.1, we have*

$$\text{vec}(\hat{\Lambda}' - \Lambda') = \mathbb{B}_1 \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} (m_i \otimes f_t) e_{it} - \mathbb{B}_2 \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^4} (m_i \otimes m_i) (e_{it}^2 - \sigma_i^2)$$

$$+ \frac{1}{T} \Delta + O_p\left(\frac{1}{N\sqrt{T}}\right) + O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T^{3/2}}\right), \quad (4.1)$$

where the symbols \mathbb{B}_1 , \mathbb{B}_2 and Δ are defined above Theorem 4.2.

The first two terms on the right hand side of (4.1) are $O_p\left(\frac{1}{\sqrt{NT}}\right)$ since their variances are $O\left(\frac{1}{NT}\right)$ and the third term is $O\left(\frac{1}{T}\right)$. The first three terms dominates the remaining terms. Theorem 4.2 reaffirms the convergence rates asserted in Theorem 4.1 and sharpens the results by explicitly giving the concrete expressions of the $O_p\left(\frac{1}{\sqrt{NT}}\right)$ and $O_p\left(\frac{1}{T}\right)$ terms. Given Theorem 4.2, invoking a Central Limit Theorem, we have the following theorem.

Theorem 4.3 (Limiting distribution) Under assumptions of Theorem 4.1, as $N, T \rightarrow \infty, N/T^2 \rightarrow 0$, we have

$$\sqrt{NT} \left[\text{vec}(\hat{\Lambda}' - \Lambda') - \frac{1}{T} \Delta \right] \xrightarrow{d} N(0, \Omega),$$

where $\Omega = \lim_{N \rightarrow \infty} \Omega_N$ with

$$\Omega_N = \mathbb{B}_1(R \otimes I_r) \mathbb{B}'_1 + \mathbb{B}_2 \left[\frac{1}{N} \sum_{i=1}^N \frac{\kappa_{iA} - \sigma_i^4}{\sigma_i^8} (m_i m'_i) \otimes (m_i m'_i) \right] \mathbb{B}'_2.$$

Theorem 4.3 shows that the MLE $\hat{\Lambda}$ has a non-negligible bias. This is in contrast to a result of Bai and Li (2012) who show that, in a high-dimensional standard factor model, the MLE is asymptotically centered around zero. Another interesting result is that the limiting variance of the MLE $\hat{\Lambda}$ depends on the kurtosis of e_{jt} . Given Theorem 4.3, when e_{it} is normally distributed, we have $\kappa_{iA} = 3\sigma_i^4$, the asymptotic variance can be simplified as the next corollary shows.

Corollary 4.1 Under assumptions of Theorem 4.3, with normality of e_{it} , we have

$$\sqrt{NT} \left[\text{vec}(\hat{\Lambda}' - \Lambda') - \frac{1}{NT} \mathbb{B}_2 \sum_{i=1}^N \frac{1}{\sigma_i^2} (m_i \otimes m_i) \right] \xrightarrow{d} N \left(0, \mathbb{B}_{1,\infty} (R_\infty \otimes I_r) \mathbb{B}'_{1,\infty} + 2\mathbb{B}_{2,\infty} V_\infty \mathbb{B}'_{2,\infty} \right),$$

where R_∞ and V_∞ are defined in Assumption C.3, $\mathbb{B}_{1,\infty}$ and $\mathbb{B}_{2,\infty}$ are almost the same as \mathbb{B}_1 and \mathbb{B}_2 except that P and R are replaced by P_∞ and R_∞ . Furthermore, if $N/T \rightarrow 0$, we have

$$\sqrt{NT} \text{vec}(\hat{\Lambda}' - \Lambda') \xrightarrow{d} N \left(0, \mathbb{B}_{1,\infty} (R_\infty \otimes I_r) \mathbb{B}'_{1,\infty} + 2\mathbb{B}_{2,\infty} V_\infty \mathbb{B}'_{2,\infty} \right).$$

Remark 4.1 To estimate the bias and the limiting variance, we use some plug-in methods. Specifically, the bias is estimated by

$$\hat{\Delta} = \hat{\mathbb{B}}_2 \frac{1}{N} \sum_{i=1}^N \frac{1}{\hat{\sigma}_i^6} (\hat{\kappa}_{iA} - \hat{\sigma}_i^4) (m_i \otimes m_i),$$

and the limiting variance is estimated by

$$\hat{\Omega} = \hat{\mathbb{B}}_1 (\hat{R} \otimes I_r) \hat{\mathbb{B}}'_1 + \hat{\mathbb{B}}_2 \left[\frac{1}{N} \sum_{i=1}^N \frac{\hat{\kappa}_{iA} - \hat{\sigma}_i^4}{\hat{\sigma}_i^8} (m_i m'_i) \otimes (m_i m'_i) \right] \hat{\mathbb{B}}'_2,$$

where

$$\begin{aligned} \hat{\mathbb{B}}_1 &= K_{kr} [(\hat{P}^{-1} \hat{\Lambda}') \otimes \hat{\Lambda}] + \hat{R}^{-1} \otimes I_r - K_{kr} (I_r \otimes \hat{\Lambda}) \hat{\mathbb{D}}_1^{-1} \mathbb{D}_2 [(\hat{P}^{-1} \hat{\Lambda}') \otimes I_r], \\ \hat{\mathbb{B}}_2 &= K_{kr} (I_r \otimes \hat{\Lambda}) \hat{\mathbb{D}}^{-1} \mathbb{D}_3 (\hat{\Lambda} \otimes \hat{\Lambda})'. \end{aligned}$$

Here $\hat{\Lambda}$ and $\hat{\sigma}_i^2$ are the MLE; $\hat{R} = \frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M$ and $\hat{P} = \frac{1}{N} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda}$; $\hat{\mathbb{D}}_1$ is almost the same as \mathbb{D}_1 except that P is replaced by \hat{P} ; $\hat{\kappa}_{iA} = \frac{1}{T} \sum_{t=1}^T \hat{e}_{it}^4$ with $\hat{e}_{it} = z_{it} - m'_i \hat{\Lambda} \hat{f}_t$ and $\hat{f}_t = (\hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} z_t$.

Remark 4.2 Theorem 4.3 is derived under a full identification of loading matrix Λ . An alternative approach to investigate the asymptotics, as adopted in Bai (2003), is that one only imposes the condition $M_{ff} = I_r$. Since in this case the original identification conditions (IC) are not met, the loading matrix Λ is not fully identified. But one can still deliver the asymptotic theory based on $\hat{\Lambda}' - \mathcal{R}\Lambda'$, where \mathcal{R} is a rotational matrix. According to (A.18) in Appendix A, together with Lemma B.3 (e), (f) and Lemma B.5 (a), we have

$$\hat{\Lambda}' - \mathcal{R}\Lambda' = \mathcal{R} \frac{1}{T} \sum_{t=1}^T f_t e_t' \Sigma_{ee}^{-1} M R_N^{-1} + O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{N\sqrt{T}}\right) + O_p\left(\frac{1}{T^{3/2}}\right),$$

where \mathcal{R} is the rotational matrix defined by

$$\mathcal{R} = \hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \Lambda + \hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T e_t f_t'$$

with $\hat{P}_N = \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda}$.

Given the above result, we have that under $N, T \rightarrow \infty, N/T^2 \rightarrow 0$,

$$\sqrt{NT} \text{vec}(\hat{\Lambda}' - \mathcal{R}\Lambda') \xrightarrow{d} N(0, R_\infty^{-1} \otimes \overline{\mathcal{R}\mathcal{R}}'),$$

where $\overline{\mathcal{R}} = \text{plim}_{N, T \rightarrow \infty} \mathcal{R}$.

Theorem 4.4 Under Assumptions A-D, as $N, T \rightarrow \infty$, we have

$$\sqrt{T}(\hat{\sigma}_i^2 - \sigma_i^2) = \frac{1}{\sqrt{T}} \sum_{t=1}^T (e_{it}^2 - \sigma_i^2) + o_p(1).$$

Given this result, we have

$$\sqrt{T}(\hat{\sigma}_i^2 - \sigma_i^2) \xrightarrow{d} N(0, \kappa_{i,4} - \sigma_i^4),$$

where $\kappa_{i,4} = E(e_{it}^4)$ is the kurtosis of e_{it} .

We emphasize that the limiting result for $\hat{\sigma}_i^2$ is independent with the identification conditions. In addition, the above limiting result is the same as that in a standard high-dimensional factor model (see, e.g., Theorem 5.4 of Bai and Li (2012)).

We finally consider the estimation of factors. Following Bai and Li (2012), we estimate the factors by the generalized least squares (GLS) method. More specifically, the GLS estimator of f_t is

$$\hat{f}_t = (\hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} z_t,$$

where $\hat{\Lambda}$ and $\hat{\Sigma}_{ee}$ are the respective MLEs of Λ and Σ_{ee} . The asymptotic representation and limiting distribution of \hat{f}_t are provided in the following theorem.

Theorem 4.5 *Under assumptions of Theorem 4.1, we have*

$$\hat{f}_t - f_t = P^{-1} \frac{1}{N} \Lambda' M' \Sigma_{ee}^{-1} e_t + O_p \left(\frac{1}{\sqrt{NT}} \right) + O_p \left(\frac{1}{T} \right),$$

where $P = \frac{1}{N} \Lambda' M' \Sigma_{ee}^{-1} M \Lambda$. Then as $N, T \rightarrow \infty$ and $N/T^2 \rightarrow 0$, we have

$$\sqrt{N}(\hat{f}_t - f_t) \xrightarrow{d} N(0, P_\infty^{-1}),$$

where $P_\infty = \lim_{N \rightarrow \infty} P$ is defined in Assumption C.3.

The above theorem indicates that the asymptotic properties of the GLS estimator for factors in the current model are the same as that in standard high-dimensional factor models^②. However, the derivation of the above theorem is actually easier due to the faster convergence rate of estimated loadings.

5 Testing

The limiting distribution of the MLE in Theorem 4.3 allows one to test whether the loading matrix Λ is equal to some known matrix. Consider the following hypothesis:

$$H_{\Lambda,0} : \Lambda = \Lambda^o, \quad H_{\Lambda,1} : \Lambda \neq \Lambda^o.$$

A Wald statistic for this hypothesis testing is

$$W_\Lambda = NT \left[\text{vec}(\hat{\Lambda}' - \Lambda^{o'}) - \frac{1}{T} \hat{\Delta} \right]' \hat{\Omega}^{-1} \left[\text{vec}(\hat{\Lambda}' - \Lambda^{o'}) - \frac{1}{T} \hat{\Delta} \right],$$

where the symbols $\hat{\Delta}$ and $\hat{\Omega}$ are given in Remark 4.1. The following theorem, which is a direct result of Theorem 4.3, gives the limiting distribution of W_Λ .

Theorem 5.1 *Under Assumptions A-D, together with IC, as $N, T \rightarrow \infty$ and $N/T^2 \rightarrow 0$, under $H_{\Lambda,0}$, we have*

$$W_\Lambda \xrightarrow{d} \chi_{kr}^2,$$

where χ_{kr}^2 denotes a chi-square distribution with degrees of freedom equal to kr .

An important issue related with the constrained factor model is that whether specification (3.1) is appropriate in a general factor model. Therefore, in practice one is likely to be interested in testing the correctness of the decomposition of loadings matrix $L = M\Lambda$. For a given M , the corresponding null and alternative hypotheses are

$$H_0 : L = M\Lambda \quad \text{for some } \Lambda,$$

^②For the asymptotic results of the GLS estimator in standard high dimensional factor models, see Theorem 6.1 of Bai and Li (2012).

$$H_1 : L \neq M\Lambda \text{ for all } \Lambda.$$

In traditional (low-dimensional) factor analysis, testing restrictions on loadings can be conducted by using the likelihood ratio (LR) principle. Because the number of parameters is finite, the number of imposed restrictions is finite too. By standard arguments, one can show that, under the null hypothesis, the LR statistic has an asymptotic χ^2 distribution with the degrees of freedom equal to the number of restrictions. In the high-dimensional setting, the number of parameters increases with the sample size. The number of restrictions possibly increases with the sample size as well. This is the case in our specification test in constrained factor models. As can be seen that under H_0 , the number of restrictions for $L = M\Lambda$ is $(N - k)r$, which proportionally increases with the number of cross sectional units. As a result, the limiting distribution of the traditional LR test would have divergent degrees of freedom, an undesirable feature which can make the test unstable. This motivates us to design a new test independent of N .

To gain an insight of our test, notice that the estimator $M\hat{\Lambda}$ ^③ under IC and H_0 should be very close to \hat{L} , the MLE of L from a standard factor model ($z_t = Lf_t + e_t$) under the identification condition that $M_{ff} = I_r$ and $\frac{1}{N}L'\Sigma_{ee}^{-1}L$ is diagonal. However, under H_1 , the two estimates will not be close to each other. Based on the above analysis, we construct the following test statistic

$$W = \sqrt{NT^2} \text{tr} \left[\frac{1}{N} (M\hat{\Lambda} - \hat{L})' \tilde{\Sigma}_{ee}^{-1} (M\hat{\Lambda} - \hat{L}) - \frac{1}{T} I_r \right],$$

where $\tilde{\Sigma}_{ee}$ is an estimator of Σ_{ee} under the alternative hypothesis.

Theorem 5.2 *Under the same assumptions of Proposition 4.1 and $N/T^2 \rightarrow 0$, under H_0 , we have*

$$W \xrightarrow{d} N(0, 2r).$$

Remark 5.1 As pointed out in Section 2, the identification condition has a sign problem. This problem should be carefully treated in the two statistics (W_Λ and W) in implementations, otherwise it may lead to an erroneous rejection of the null hypothesis. To eliminate this problem, when calculating W_Λ , we first compute the inter product of each column of $\hat{\Lambda}$ and the counterpart of Λ^o . If the value is negative, we multiple -1 on this column of $\hat{\Lambda}$. As regard to W , both \hat{L} and $M\hat{\Lambda}$ have the sign problem, but we can use a similar procedure to deal with it. That is, for each column of \hat{L} , we calculate the inner product of this column and its counterpart of $M\hat{\Lambda}$. If the inner product is negative, we multiple -1 on this column of \hat{L} . After this treatment, the sign problem concomitant with the identification condition is removed.

^③An alternative estimator is $M\hat{\Lambda}^\dagger$, where $\hat{\Lambda}^\dagger$ is the bias-corrected estimator for Λ . It can be shown that the difference of the two statistics (which are based on $\hat{\Lambda}^\dagger$ and $\hat{\Lambda}$) is asymptotically negligible under $N/T^2 \rightarrow 0$.

Remark 5.2 Although we use the symbol W to denote the proposed statistic in the paper, our W statistic differs from the conventional Wald test. There are some key features that are different between our W test and the Wald test. First, the Wald test only involves estimators from an unconstrained model. In contrast, we use estimators from both constrained and unconstrained models to construct the W statistic. Second, the Wald test has an asymptotic χ^2 distribution with the value of degrees of freedom equal to the number of restrictions. But our W statistic has an asymptotic normal distribution, which is free of degree of freedom. For the same reasons, our W statistic is also different from a conventional Lagrange multiplier test.

6 Partially Constrained Factor Models

In this section, we consider the following partially constrained factor model

$$z_t = M\Lambda f_t + \Gamma g_t + e_t \triangleq \Phi h_t + e_t, \quad (6.1)$$

where $\Phi = [M\Lambda, \Gamma]$, $h_t = (f_t', g_t')$ is an r -dimensional vector, f_t is an r_1 -dimensional vector and g_t an r_2 -dimensional vector with $r_1 + r_2 = r$. Again we study the ML estimation on model (6.1).

To analyze the MLE, we make the following assumptions.

Assumption A'. The factors $\{h_t\}$ satisfy the conditions in Assumption A.

Assumption C'. There exists a positive constant C such that $\|\phi_i\| < C$ for all i , where ϕ_i is the transpose of the i th row of Φ . Let $\mathcal{H} = \frac{1}{N}\Phi'\Sigma_{ee}^{-1}\Phi$, we assume $\bar{\mathcal{H}} = \lim_{N \rightarrow \infty} \mathcal{H} > 0$.

Identification condition, IC'. The identification conditions considered here are similar to those in the pure constrained factor model. More specifically, we require that $M_{hh} = \frac{1}{T}\sum_{t=1}^T h_t h_t' = I_r$ and \mathcal{H} is a diagonal matrix with all its diagonal elements distinct and arranged in a descending order.

Let $\Sigma_{zz} = \Phi\Phi' + \Sigma_{ee}$ and $\theta = (\Lambda, \Gamma, \Sigma_{ee})$. The MLE is defined as

$$\hat{\theta} = \operatorname{argmax}_{\theta \in \Theta} \mathcal{L}(\theta),$$

where

$$\mathcal{L}(\theta) = -\frac{1}{2N} \ln |\Sigma_{zz}| - \frac{1}{2N} \operatorname{tr}[M_{zz}\Sigma_{zz}^{-1}].$$

Here Θ is the parameter space specified by Assumption D and the identification condition IC'. In the supplementary appendix D (available upon request), we show that the first order condition for Λ can be written as

$$\hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} (M_{zz} - \hat{\Sigma}_{zz}) \hat{\Sigma}_{ee}^{-1} M = 0. \quad (6.2)$$

The first order condition for Γ can be written as

$$\hat{\Gamma}' \hat{\Sigma}_{ee}^{-1} (M_{zz} - \hat{\Sigma}_{zz}) = 0. \quad (6.3)$$

The first order condition for Σ_{ee} can be written as

$$\text{diag} \left[(M_{zz} - \hat{\Sigma}_{zz}) - M \hat{\Lambda} \hat{G}_1 \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} (M_{zz} - \hat{\Sigma}_{zz}) - (M_{zz} - \hat{\Sigma}_{zz}) \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{G}_1 \hat{\Lambda}' M' \right] = 0. \quad (6.4)$$

Before we present the asymptotic results for the MLE, we first introduce some notation

$$\begin{aligned} \mathbb{B}_1^* &= R^{-1} \otimes I_{r_1} + K_{kr_1} [(P^{-1} \Lambda') \otimes \Lambda] - K_{kr_1} (E_1' \otimes \Psi) \mathbb{D}_1^{-1} \mathbb{D}_2 [(\mathcal{H}^{-1} E_1 \Lambda') \otimes E_1], \\ \mathbb{B}_2^* &= K_{kr_1} [P^{-1} \otimes \psi] - K_{kr_1} (E_1' \otimes \Psi) \mathbb{D}_1^{-1} \mathbb{D}_2 [(\mathcal{H}^{-1} E_1) \otimes E_2], \\ \mathbb{B}_3^* &= -K_{kr_1} (E_1' \otimes \Psi) \mathbb{D}_1^{-1} \mathbb{D}_2 [(\mathcal{H}^{-1} E_2) \otimes E_1], \\ \mathbb{B}_4^* &= -K_{kr_1} (E_1' \otimes \Psi) \mathbb{D}_1^{-1} \mathbb{D}_2 [(\mathcal{H}^{-1} E_2) \otimes E_2], \quad \mathbb{B}_5^* = -K_{kr_1} (E_1' \otimes \Psi) \mathbb{D}_1^{-1} \mathbb{D}_3, \\ \Delta^* &= K_{kr_1} (E_1' \otimes \Psi) \mathbb{D}_1^{-1} \mathbb{D}_3 \left[\frac{1}{N} \sum_{i=1}^N \frac{1}{\sigma_i^6} (\phi_i \otimes \phi_i) (\kappa_{i,4} - \sigma_i^4) + \text{vec}(r_1 \mathcal{H} - E_2 E_2') \right], \end{aligned}$$

where $E_1 = [I_{r_1}, 0_{r_1 \times r_2}]'$, $E_2 = [0_{r_2 \times r_1}, I_{r_2}]'$, $\psi = (M' \Sigma_{ee}^{-1} M)^{-1} M' \Sigma_{ee}^{-1} \Gamma$, $\Psi = [\Lambda, \psi]$ and \mathcal{H} is defined in Assumption C'. The symbols $\kappa_{i,4}$, K_{mm} , P , R , \mathbb{D}_1 , \mathbb{D}_2 and \mathbb{D}_3 are defined the same as in Section 4.

Let γ_i be the transpose of the i th row of Γ . The following theorem states the asymptotic representations for the MLE. The consistency and convergence rates are implied by the theorem.

Theorem 6.1 *Under Assumptions A', B, C' and D, when $N, T \rightarrow \infty$, we have, for all i ,*

$$\hat{\sigma}_i^2 - \sigma_i^2 = \frac{1}{T} \sum_{t=1}^T (e_{it}^2 - \sigma_i^2) + O_p \left(\frac{1}{T} \right).$$

In addition, if IC' is imposed, we have, for all i ,

$$\hat{\gamma}_i - \gamma_i = \frac{1}{T} \sum_{t=1}^T g_t e_{it} + O_p \left(\frac{1}{T} \right)$$

and

$$\begin{aligned} \text{vec}(\hat{\Lambda}' - \Lambda') &= \mathbb{B}_1^* \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} (m_i \otimes f_t) e_{it} + \mathbb{B}_2^* \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} (\Lambda' m_i \otimes g_t) e_{it} \\ &+ \mathbb{B}_3^* \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} (\gamma_i \otimes f_t) e_{it} + \mathbb{B}_4^* \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} (\gamma_i \otimes g_t) e_{it} \\ &+ \mathbb{B}_5^* \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^4} (\phi_i \otimes \phi_i) (e_{it}^2 - \sigma_i^2) + \frac{1}{T} \Delta^* \\ &+ O_p \left(\frac{1}{N\sqrt{T}} \right) + O_p \left(\frac{1}{\sqrt{NT}} \right) + O_p \left(\frac{1}{T^{3/2}} \right), \end{aligned}$$

where $\mathbb{B}_1^*, \dots, \mathbb{B}_5^*$ and Δ^* are defined above this theorem.

Given the above theorem, we have the following distribution results for the MLE.

Corollary 6.1 *Under Assumptions A', B, C' and D, when $N, T \rightarrow \infty$, we have, for all i ,*

$$\sqrt{T}(\hat{\sigma}_i^2 - \sigma_i^2) \xrightarrow{d} N(0, \kappa_{i,4} - \sigma_i^4).$$

In addition, if IC' is imposed, we have, for all i ,

$$\sqrt{T}(\hat{\gamma}_i - \gamma_i) \xrightarrow{d} N(0, \sigma_i^2 I_{r_2}).$$

If $N/T^2 \rightarrow 0$ is further imposed, we have

$$\sqrt{NT} \left[\text{vec}(\hat{\Lambda}' - \Lambda') - \frac{1}{T} \Delta^* \right] \xrightarrow{d} N(0, \Omega^*),$$

where $\Omega^ = \lim_{N \rightarrow \infty} \Omega_N^*$ with*

$$\begin{aligned} \Omega_N^* &= \mathbb{B}_1^*(R \otimes I_{r_1}) \mathbb{B}_1^{*'} + \mathbb{B}_2^*(P \otimes I_{r_1}) \mathbb{B}_2^{*'} + \mathbb{B}_3^*(Q \otimes I_{r_1}) \mathbb{B}_3^{*'} + \mathbb{B}_4^*(Q \otimes I_{r_2}) \mathbb{B}_4^{*'} \\ &\quad + \mathbb{B}_1^*(S \otimes I_{r_1}) \mathbb{B}_3^{*'} + \mathbb{B}_3^*(S' \otimes I_{r_1}) \mathbb{B}_1^{*'} + \mathbb{B}_5^* \left[\frac{1}{N} \sum_{i=1}^N \frac{1}{\sigma_i^8} (\phi_i \phi_i') \otimes (\phi_i \phi_i') (\kappa_{i,4} - \sigma_i^4) \right] \mathbb{B}_5^{*'}, \end{aligned}$$

where $Q = \Gamma' \Sigma_{ee}^{-1} \Gamma / N$ and $S = M' \Sigma_{ee}^{-1} \Gamma / N$.

The approach to estimate the factors in partially constrained factor models is similar as before. Given the MLE $\hat{\Lambda}, \hat{\Gamma}$ and $\hat{\Sigma}_{ee}$, the GLS estimator of h_t is

$$\hat{h}_t = (\hat{\Phi}' \hat{\Sigma}_{ee}^{-1} \hat{\Phi})^{-1} \hat{\Phi}' \hat{\Sigma}_{ee}^{-1} z_t,$$

where $\hat{\Phi} = (M\hat{\Lambda}, \hat{\Gamma})$. Using the similar arguments as in the proof of Theorem 4.5, we have the following asymptotic representation and limiting distribution results on \hat{h}_t .

Theorem 6.2 *Under Assumptions A', B, C' and D, together with IC', we have, for all t ,*

$$\hat{h}_t - h_t = \mathcal{H}^{-1} \frac{1}{N} \Phi' \Sigma_{ee}^{-1} e_t + O_p \left(\frac{1}{\sqrt{NT}} \right) + O_p \left(\frac{1}{T} \right),$$

where $\mathcal{H} = \frac{1}{N} \Phi' \Sigma_{ee}^{-1} \Phi$. Then as $N, T \rightarrow \infty$ and $N/T^2 \rightarrow 0$, we have

$$\sqrt{N}(\hat{h}_t - h_t) \xrightarrow{d} N(0, \bar{\mathcal{H}}^{-1}),$$

where $\bar{\mathcal{H}} = \lim_{N \rightarrow \infty} \mathcal{H}$ is defined in Assumption C'.

7 EM algorithm

The ML estimation can be easily implemented via the EM algorithm. The iterating formulas for a purely constrained factor model and a partially constrained one are different. We present them separately.

7.1 EM algorithm for the pure constrained factor model

Let $\theta^{(k)} = (\Lambda^{(k)}, \Sigma_{ee}^{(k)})$ denote the estimate at the k th iteration. The EM algorithm updates and calculates $\theta^{(k+1)} = (\Lambda^{(k+1)}, \Sigma_{ee}^{(k+1)})$ by

$$\Lambda^{(k+1)} = (M' \Sigma_{ee}^{(k)-1} M)^{-1} \left[M' \Sigma_{ee}^{(k)-1} \frac{1}{T} \sum_{t=1}^T E(z_t f_t' | Z, \theta^{(k)}) \right] \left[\frac{1}{T} \sum_{t=1}^T E(f_t f_t' | Z, \theta^{(k)}) \right]^{-1},$$

$$\text{diag}(\Sigma_{ee}^{(k+1)}) = \text{diag} \left\{ M_{zz} - \frac{2}{T} \sum_{t=1}^T E(z_t f_t' | Z, \theta^{(k)}) \Lambda^{(k+1)'} M' \right. \\ \left. + M \Lambda^{(k+1)} \frac{1}{T} \sum_{t=1}^T E(f_t f_t' | Z, \theta^{(k)}) \Lambda^{(k+1)'} M' \right\},$$

where $\Sigma_{zz}^{(k)} = M \Lambda^{(k)} \Lambda^{(k)'} M' + \Sigma_{ee}^{(k)}$ and

$$\frac{1}{T} \sum_{t=1}^T E(f_t f_t' | Z, \theta^{(k)}) = \Lambda^{(k)'} M' (\Sigma_{zz}^{(k)})^{-1} M_{zz} (\Sigma_{zz}^{(k)})^{-1} M \Lambda^{(k)} + I_r - \Lambda^{(k)'} M' (\Sigma_{zz}^{(k)})^{-1} M \Lambda^{(k)},$$

$$\frac{1}{T} \sum_{t=1}^T E(z_t f_t' | Z, \theta^{(k)}) = M_{zz} (\Sigma_{zz}^{(k)})^{-1} M \Lambda^{(k)}.$$

The above iteration continues until $\|\theta^{(k+1)} - \theta^{(k)}\|$ is smaller than a preset tolerance. For the initial values, the PC estimates proposed in Tsai and Tsay (2010) are recommended. When iterations are terminated, the estimates, denoted by $(\Lambda^+, \Sigma_{ee}^+)$, need to be further normalized to satisfy the identification conditions in Section 3. The normalization can be conducted as follows. Let V^+ be the orthogonal matrix consisting of the eigenvectors of the matrix $\frac{1}{N} \Lambda^+ M' (\Sigma_{ee}^+)^{-1} M \Lambda^+$ with the corresponding eigenvalues arranged in a descending order. Let $\hat{\Lambda} = \Lambda^+ V^+$ and $\hat{\Sigma}_{ee} = \Sigma_{ee}^+$. Then $\hat{\theta} = (\hat{\Lambda}, \hat{\Sigma}_{ee})$ is the MLE that satisfies IC.

Bai and Li (2012) show that the iterating formulas of the EM algorithm approach to the first order conditions of the likelihood function as the iteration tends to infinity. Using their arguments, one can show similar results in constrained factor models. Since the proof is almost the same as in Bai and Li (2012), we omit it for sake of space.

7.2 EM algorithm for the partially constrained factor model

Let $\theta^{(k)} = (\Lambda^{(k)}, \Gamma^{(k)}, \Sigma_{ee}^{(k)})$ denote the estimate at the k th iteration. The EM algorithm updates and calculates $\theta^{(k+1)} = (\Lambda^{(k+1)}, \Gamma^{(k+1)}, \Sigma_{ee}^{(k+1)})$ by

$$\Lambda^{(k+1)} = (M' \Sigma_{ee}^{(k)-1} M)^{-1} \left[M' \Sigma_{ee}^{(k)-1} \frac{1}{T} \sum_{t=1}^T E(z_t f_t' | Z, \theta^{(k)}) \right] \left[\frac{1}{T} \sum_{t=1}^T E(f_t f_t' | Z, \theta^{(k)}) \right]^{-1}$$

$$- (M' \Sigma_{ee}^{(k)-1} M)^{-1} \left[M' \Sigma_{ee}^{(k)-1} \Gamma^{(k)} \frac{1}{T} \sum_{t=1}^T E(g_t f_t' | Z, \theta^{(k)}) \right] \left[\frac{1}{T} \sum_{t=1}^T E(f_t f_t' | Z, \theta^{(k)}) \right]^{-1},$$

$$\begin{aligned}
\Gamma^{(k+1)} &= \left[\frac{1}{T} \sum_{t=1}^T E(z_t g_t' | Z, \theta^{(k)}) \right] \left[\frac{1}{T} \sum_{t=1}^T E(g_t g_t' | Z, \theta^{(k)}) \right]^{-1} \\
&\quad - \left[M\Lambda^{(k+1)} \frac{1}{T} \sum_{t=1}^T E(f_t g_t' | Z, \theta^{(k)}) \right] \left[\frac{1}{T} \sum_{t=1}^T E(g_t g_t' | Z, \theta^{(k)}) \right]^{-1}, \\
\text{diag}(\Sigma_{ee}^{(k+1)}) &= \text{diag} \left\{ M_{zz} - \frac{2}{T} \sum_{t=1}^T E(z_t f_t' | Z, \theta^{(k)}) \Lambda^{(k+1)'} M' - \frac{2}{T} \sum_{t=1}^T E(z_t g_t' | Z, \theta^{(k)}) \Gamma^{(k+1)'} \right. \\
&\quad + M\Lambda^{(k+1)} \frac{1}{T} \sum_{t=1}^T E(f_t f_t' | Z, \theta^{(k)}) \Lambda^{(k+1)'} M' + \Gamma^{(k+1)} \frac{1}{T} \sum_{t=1}^T E(g_t g_t' | Z, \theta^{(k)}) \Gamma^{(k+1)'} \\
&\quad \left. + 2M\Lambda^{(k+1)} \frac{1}{T} \sum_{t=1}^T E(f_t g_t' | Z, \theta^{(k)}) \Gamma^{(k+1)'} \right\},
\end{aligned}$$

where $\Sigma_{zz}^{(k)} = M\Lambda^{(k)} \Lambda^{(k)'} M' + \Gamma^{(k)} \Gamma^{(k)'} + \Sigma_{ee}^{(k)}$ and

$$\begin{aligned}
\frac{1}{T} \sum_{t=1}^T E(f_t f_t' | Z, \theta^{(k)}) &= \Lambda^{(k)'} M' (\Sigma_{zz}^{(k)})^{-1} M_{zz} (\Sigma_{zz}^{(k)})^{-1} M\Lambda^{(k)} + I_{r_1} - \Lambda^{(k)'} M' (\Sigma_{zz}^{(k)})^{-1} M\Lambda^{(k)}, \\
\frac{1}{T} \sum_{t=1}^T E(f_t g_t' | Z, \theta^{(k)}) &= \Lambda^{(k)'} M' (\Sigma_{zz}^{(k)})^{-1} M_{zz} (\Sigma_{zz}^{(k)})^{-1} \Gamma^{(k)} - \Lambda^{(k)'} M' (\Sigma_{zz}^{(k)})^{-1} \Gamma^{(k)}, \\
\frac{1}{T} \sum_{t=1}^T E(g_t g_t' | Z, \theta^{(k)}) &= \Gamma^{(k)'} (\Sigma_{zz}^{(k)})^{-1} M_{zz} (\Sigma_{zz}^{(k)})^{-1} \Gamma^{(k)} + I_{r_2} - \Gamma^{(k)'} (\Sigma_{zz}^{(k)})^{-1} \Gamma^{(k)}, \\
\frac{1}{T} \sum_{t=1}^T E(z_t f_t' | Z, \theta^{(k)}) &= M_{zz} (\Sigma_{zz}^{(k)})^{-1} M\Lambda^{(k)}, \\
\frac{1}{T} \sum_{t=1}^T E(z_t g_t' | Z, \theta^{(k)}) &= M_{zz} (\Sigma_{zz}^{(k)})^{-1} \Gamma^{(k)}.
\end{aligned}$$

Likewise, we use the PC estimates as the starting values, and iterate the above formulas until $\|\theta^{(k+1)} - \theta^{(k)}\|$ is smaller than a preset tolerance. Let $\theta^\diamond = (\Lambda^\diamond, \Gamma^\diamond, \Sigma_{ee}^\diamond)$ be the estimates of the last iteration. Again we need rotate θ^\diamond to satisfy the IC''. Let V^\diamond be the orthogonal matrix consisting of the eigenvectors of the matrix $\frac{1}{N} \Phi^{\diamond'} (\Sigma_{ee}^\diamond)^{-1} \Phi^\diamond$ with the corresponding eigenvalues arranged in a descending order, where $\Phi^\diamond = (M\Lambda^\diamond, \Gamma^\diamond)$. Let $\Phi^\diamond V^\diamond$ and split Φ^\diamond into $\Phi^\Delta = (\Phi_1^\Delta, \Phi_2^\Delta)$, where Φ_1^Δ is made up with the left r_1 columns and Φ_2^Δ the remaining r_2 columns. Then calculate $\hat{\Lambda} = (M'M)^{-1} M' \Phi_1^\Delta$, and simply let $\hat{\Gamma} = \Phi_2^\Delta$ and $\hat{\Sigma}_{ee} = \Sigma_{ee}^\diamond$. Then $\hat{\theta} = (\hat{\Lambda}, \hat{\Gamma}, \hat{\Sigma}_{ee})$ is the MLE that satisfies IC''.

Again, we can show that the limit of the iterated EM solutions satisfy the first order conditions (6.2), (6.3) and (6.4). The proof is similar to the pure constrained factor model case and therefore skipped here.

8 Simulation results

In this section, we run simulations to investigate the finite sample performance of the MLE, as well as the empirical size and power of the W test.

8.1 Finite sample performance of the MLE

We first conduct simulations to investigate the finite sample properties of the MLE and compare it with the PC estimates proposed by Tsai and Tsay (2010).

In the literature on high dimensional factor models, researchers usually use a generalized R^2 or a trace ratio to measure the goodness-of-fit, e.g., Stock and Watson (2002), Doz, Giannone and Reichlin (2012) and Bai and Li (2012). These measures are invariant to the rotational indeterminacy and therefore effective to perform the measure task. However, in constrained factor models, such measures are not suitable since the estimates have faster convergence rates, which often leads to a high value of the generalized R^2 or the trace ratio. For this reason, we instead consider an alternative measure by rotating the underlying values to satisfy the identification condition and investigating the precision of $\hat{\Lambda} - \Lambda$ for rotated values. We calculate the mean absolute deviation (MAD) and the root mean square error (RMSE) based on the rotated underlying values. We also calculate the root asymptotic variance (RAvar) to check the convergence rate of $\hat{\Lambda}$ presented in Theorem 4.1. The calculation formulas based on S simulations are as follows

$$\begin{aligned} \text{MAD} &= \frac{1}{S} \sum_{s=1}^S \left(\frac{1}{kr} \sum_{p=1}^k \sum_{i=1}^r |\hat{\Lambda}_{pi}^s - \Lambda_{pi}^s| \right), \\ \text{RMSE} &= \sqrt{\frac{1}{S} \sum_{s=1}^S \left(\frac{1}{kr} \sum_{p=1}^k \sum_{i=1}^r (\hat{\Lambda}_{pi}^s - \Lambda_{pi}^s)^2 \right)}, \\ \text{RAvar} &= \sqrt{NT} \times \sqrt{\frac{1}{S} \sum_{s=1}^S \left(\frac{1}{kr} \sum_{p=1}^k \sum_{i=1}^r (\hat{\Lambda}_{pi}^{bc,s} - \Lambda_{pi}^s)^2 \right)}, \end{aligned}$$

where $\hat{\Lambda}_{pi}^s$ and $\hat{\Lambda}_{pi}^{bc,s}$ are the MLE and biased-corrected MLE in the s th simulation, respectively.

Data are generated according to $z_t = M\Lambda f_t + e_t$, where all elements of M are drawn independently from $U[0, 1]$ and all elements of Λ and F independently from $N(0, 1)$. The idiosyncratic errors e_{it} are generated according to $e_{it} = \sigma_i \epsilon_{it}$ with σ_i^2 being the i th diagonal element of $(M\Lambda\Lambda'M')$ multiplying $\frac{b_i}{1-b_i}$, where $b_i = 0.2 + 0.6U_i$ and $U_i \sim U[0, 1]$. The component ϵ_{it} is generated from the three distributions: the normal distribution, student's distribution with 5 degrees of freedom and chi-squared distribution with 2 degrees of freedom. For the latter two distributions, we normalize the random variable to have zero mean and unit variance. For the values of k and r , we consider two cases: $(k, r) = (3, 1)$ and $(k, r) = (8, 3)$.

Throughout this section, we assume that the number of common factors is known. There are a number of methods at hand to determine the number of factors, for example, the information criterion method by Bai and Ng (2002), the largest eigenvalue-ratios method by Ahn and Horenstein (2013) and the eigenvalue empirical distribution method by Onatski (2010). If the number of factors is unknown, one can choose any of the

method mentioned above to estimate it. Tables 1 and 2 present the performance of the MLE and the PC estimate for normal errors under the sample sizes of $N = 30, 50, 100, 150$ and $T = 30, 50, 100$. The results under student-t errors and chi-square errors are almost the same as those for normal errors and are given in Table E1-E4 in the supplementary appendix E (available upon request) to save space. All these results are obtained based on 1000 repetitions.

From Tables 1 and 2, we can see that both MAD and RMSE of the MLE are much smaller than those of PC estimates for all (N, T) combinations, implying that the MLE performs better than the PC estimate. Regarding the RAvar^④, we see that the MLE has almost constant RAvars when the time dimension T or the cross section dimension N increases, implying that the convergence rate of the MLE is \sqrt{NT} . This simulation result is consistent with our theoretical results in Section 4. In addition, it seems that the PC estimate also has \sqrt{NT} convergence rate from simulations. Finally, we note that the RMSEs of the MLE are smaller than those of the PC estimates, indicating that the MLE is more efficient than the PC estimates.

Table 1: $k = 3, r = 1$, and $\epsilon_{it} \sim N(0, 1)$.

$\Lambda_{3 \times 1}$		MLE			PC		
N	T	MAD	RMSE	RAvar	MAD	RMSE	RAvar
30	30	0.0440	0.0716	2.2301	0.0943	0.1386	N/A
50	30	0.0349	0.0540	1.9887	0.0654	0.0934	N/A
100	30	0.0262	0.0417	2.0504	0.0474	0.0677	N/A
150	30	0.0216	0.0340	2.1741	0.0410	0.0582	N/A
30	50	0.0333	0.0533	2.1936	0.0787	0.1145	N/A
50	50	0.0237	0.0368	1.9426	0.0546	0.0800	N/A
100	50	0.0190	0.0306	1.9194	0.0375	0.0541	N/A
150	50	0.0159	0.0255	2.0863	0.0293	0.0417	N/A
30	100	0.0232	0.0374	2.1425	0.0674	0.0964	N/A
50	100	0.0172	0.0263	1.8314	0.0443	0.0611	N/A
100	100	0.0105	0.0168	1.7473	0.0253	0.0358	N/A
150	100	0.0102	0.0165	1.8668	0.0200	0.0288	N/A

Table 2: $k = 8, r = 3$, and $\epsilon_{it} \sim N(0, 1)$.

^④Since we do not know whether the PC estimate is biased, and if biased, what is the bias formula. Hence, we cannot calculate RAvar for the PC estimate.

$\Lambda_{8 \times 3}$		MLE			PC		
N	T	MAD	RMSE	RAvar	MAD	RMSE	RAvar
30	30	0.3498	0.5006	15.2632	0.5655	0.8071	N/A
50	30	0.2307	0.3310	13.6988	0.3744	0.5363	N/A
100	30	0.1537	0.2307	12.5998	0.2224	0.3131	N/A
150	30	0.1245	0.1881	11.7159	0.1735	0.2452	N/A
30	50	0.2637	0.3744	14.4701	0.5130	0.7521	N/A
50	50	0.1794	0.2689	13.1269	0.3184	0.4679	N/A
100	50	0.1082	0.1578	12.1691	0.1763	0.2545	N/A
150	50	0.0860	0.1291	12.3152	0.1382	0.2091	N/A
30	100	0.1846	0.2698	15.5540	0.4570	0.6882	N/A
50	100	0.1213	0.1937	13.3273	0.2622	0.4064	N/A
100	100	0.0774	0.1258	11.9418	0.1440	0.2157	N/A
150	100	0.0620	0.1021	12.9696	0.1033	0.1633	N/A

8.2 Empirical size of the W test

In this subsection, we use simulations to study the empirical size of the W statistic. The data generating process is the same as in previous subsection, but with more combinations of (N, T) . We investigate the performance of W under three nominal levels 1%, 5% and 10%. The empirical sizes of W for the case $(k, r) = (3, 1)$ are given in Table 3, which is obtained from 1000 repetitions.

Table 3: The empirical size of the test statistic W for $(k, r) = (3, 1)$

Empirical size of W										
$\epsilon_{it} \sim$		$N(0,1)$			t_5			$\chi^2(2)$		
N	T	1%	5%	10%	1%	5%	10%	1%	5%	10%
30	30	3.6%	7.4%	13.5%	3.8%	8.5%	12.9%	2.7%	8.0%	13.3%
50	30	4.4%	11.5%	16.6%	3.9%	9.5%	16.3%	5.4%	10.5%	16.1%
100	30	6.7%	14.2%	20.5%	6.5%	13.9%	20.1%	5.5%	12.9%	21.1%
150	30	9.2%	18.4%	24.8%	8.1%	18.6%	27.1%	8.2%	20.3%	29.0%
30	50	1.7%	5.9%	11.3%	1.3%	5.8%	12.7%	1.7%	6.6%	11.6%
50	50	3.1%	6.8%	13.0%	2.6%	6.1%	11.0%	2.0%	7.0%	12.1%
100	50	3.3%	8.0%	15.2%	2.3%	8.3%	14.2%	3.5%	9.7%	15.7%
150	50	4.6%	11.4%	18.1%	3.4%	11.1%	17.3%	2.8%	9.3%	15.8%
30	100	0.6%	4.5%	10.4%	1.4%	4.0%	10.6%	1.0%	4.8%	10.9%
50	100	1.5%	4.2%	10.9%	1.5%	6.1%	9.9%	1.2%	5.8%	11.7%
100	100	1.4%	6.5%	11.6%	0.9%	5.8%	12.6%	1.5%	6.5%	12.4%
150	100	1.6%	5.6%	10.9%	2.0%	7.5%	12.7%	1.9%	5.8%	11.3%
30	150	0.6%	5.0%	10.5%	1.0%	5.0%	9.9%	1.2%	5.8%	10.2%
50	150	1.5%	5.9%	10.4%	1.5%	4.8%	10.2%	1.5%	5.1%	9.6%
100	150	0.7%	6.2%	10.7%	1.2%	5.4%	10.2%	1.5%	5.8%	11.6%
150	150	1.9%	5.9%	9.6%	1.6%	5.0%	11.5%	1.7%	5.2%	10.8%
100	100	1.4%	6.5%	11.6%	0.9%	5.8%	12.6%	1.5%	6.5%	12.4%
200	100	1.3%	6.1%	11.2%	1.4%	6.7%	13.5%	2.2%	7.2%	12.6%
300	100	2.3%	6.5%	12.8%	2.1%	6.8%	12.7%	1.8%	7.9%	12.9%
100	200	1.3%	4.0%	9.4%	1.3%	5.3%	10.8%	1.1%	5.1%	11.3%
200	200	1.4%	5.6%	10.5%	0.9%	4.9%	9.6%	1.4%	6.1%	11.6%
300	200	1.3%	6.1%	8.6%	1.5%	5.4%	11.6%	1.5%	5.9%	11.7%
100	300	0.4%	4.5%	9.5%	1.2%	5.1%	11.8%	1.2%	5.1%	9.2%
200	300	0.9%	6.1%	10.5%	1.3%	4.9%	9.1%	0.8%	6.2%	11.6%
300	300	1.3%	5.2%	10.9%	0.7%	3.9%	8.5%	1.2%	4.4%	9.0%
100	500	0.8%	5.3%	9.8%	0.8%	4.6%	10.9%	1.1%	5.2%	9.7%
200	500	0.9%	5.4%	9.8%	0.5%	5.1%	9.8%	1.0%	5.2%	10.3%
300	500	0.6%	5.3%	10.5%	1.5%	5.9%	9.2%	0.9%	5.0%	9.4%

From the results in Table 3, we emphasize the following findings. First, the performance of the W test is considerably good overall. Except for the sample size when T is small, almost all the empirical sizes of the W statistic fall in the interval $[5\%, 10\%]$ under the 5% nominal level. Second, the distribution type of errors has no significant impact on the performance of W . The W statistic performs very similarly under three different error distributions. This is consistent with the theoretical result in Section 5. Third, the performance of W is closely linked with time period number T , loosely with the number of units N . For example, when $T = 30$, the W statistic suffers a mildly severe size distortion. But when T grows to 50, the size distortion considerably decreases. As regard to N , we see that the W statistic performs well even when $N = 30$. We conjecture the reason is that when T is small, the variance σ_i^2 are estimated inaccurately, which leads to a poor performance of W .

Tsai and Tsay (2010) propose using a traditional likelihood ratio (LR) statistic to perform model specification testing. In the factor model literature, LR tests are usually considered under the fixed- N , large- T setup, see Lawley and Maxwell (1971). As mentioned in the introduction, when N and T are both large the traditional LR test may not be suitable. For example, the adjusted likelihood ratio test, which is often used with

consideration of finite sample performance, may be negative for too large N . According to the simulation results in Table 7 in Tsai and Tsay (2010), the LR test suffers size distortion issue even when N is not large. As a primary competitor to our W statistic, we compare the performance of the W statistic and the LR one under the current data generating setup. We find that the performance of the W statistic dominates that of the LR test. Details are given in Appendix F in the supplementary material of this paper.

8.3 Empirical power of the W test

We next study the empirical power of the W test. Data are generated by $z_t = Lf_t + e_t$ with

$$L = M\Lambda + d \cdot v,$$

where M, Λ, f_t and e_t are generated in the same way as in Section 8.1. The symbol v is an $N \times r$ noise matrix with its elements drawn from $N(0, 1)$ and d is a prespecified constant, which is related with N and T and is used to control the magnitude of deviation from the null hypothesis. In this section, we set it as

$$d = \frac{\alpha}{\sqrt[4]{N}\sqrt{T}}$$

with $\alpha = 0.2, 0.5, 2$ and 5 . In classical models, if an estimator is \sqrt{T} -consistent, the local power is studied under $\beta = \beta^* + \frac{1}{\sqrt{T}}\alpha$, where β^* denotes the true value. However, this general result cannot be applied to the present context since we renormalize the distance between estimators from the constrained and unconstrained models to accommodate the large number of restrictions imposed in the null hypothesis. Directly deriving the local power of W is challenging. We conjecture that the W statistic can detect local alternatives that approach the null model at a rate of $N^{-1/4}T^{-1/2}$. Simulation results below seem to support our conjecture since the local power converges to some value as N and T grow larger in all choices of α .

Table 4 presents the empirical power of the W test for the case $(k, r) = (3, 1)$ under normal errors. It is seen that the W statistic has higher power when α is larger and lower power when α is smaller. This is an expected result. As α becomes larger, the distance between the null hypothesis and the alternative hypothesis is larger and then we have more chances to differentiate the two hypotheses. Given that the W statistic has considerable power even against the local alternatives that are $N^{-1/4}T^{-1/2}$ away from the null model, we conclude that the W has good performance in terms of empirical power. We also compare empirical powers of the W statistic and the LR test. We find that the performance of the W test is better than that of the LR test. Details are given in the supplementary Appendix F.

Table 4: The empirical power of the W test for $(k, r) = (3, 1)$

α		Empirical power of W											
		0.2			0.5			2			5		
N	T	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%
30	30	22.9%	31.4%	37.4%	52.0%	57.5%	61.7%	91.2%	93.1%	93.7%	99.7%	100.0%	100.0%
50	30	31.8%	39.4%	44.9%	58.2%	64.1%	67.5%	94.1%	95.7%	96.4%	100.0%	100.0%	100.0%
100	30	51.4%	59.4%	63.7%	71.4%	77.3%	81.1%	96.2%	98.0%	98.7%	100.0%	100.0%	100.0%
150	30	55.5%	63.9%	68.0%	74.4%	78.9%	81.6%	97.9%	98.9%	99.2%	100.0%	100.0%	100.0%
30	50	22.9%	30.3%	35.2%	51.1%	57.4%	60.7%	89.3%	91.9%	93.6%	99.6%	99.8%	99.8%
50	50	29.2%	36.3%	42.2%	58.2%	63.8%	67.4%	93.7%	95.8%	96.7%	99.8%	99.9%	99.9%
100	50	45.5%	51.7%	56.3%	69.2%	72.7%	76.1%	96.5%	97.7%	98.1%	100.0%	100.0%	100.0%
150	50	51.3%	58.3%	63.4%	70.9%	76.0%	79.2%	97.3%	98.2%	98.5%	100.0%	100.0%	100.0%
30	100	20.5%	25.7%	31.5%	53.6%	60.7%	62.9%	90.0%	92.2%	93.8%	99.5%	99.6%	99.6%
50	100	29.8%	35.6%	41.1%	59.3%	64.2%	67.2%	93.1%	94.7%	95.7%	100.0%	100.0%	100.0%
100	100	37.7%	43.3%	47.5%	65.6%	70.1%	72.3%	94.1%	96.2%	97.3%	99.9%	100.0%	100.0%
150	100	49.8%	55.4%	59.0%	70.1%	74.2%	77.6%	95.5%	96.6%	97.2%	100.0%	100.0%	100.0%
30	150	19.9%	25.4%	29.8%	55.8%	62.1%	64.5%	88.2%	91.2%	92.0%	99.6%	99.8%	99.9%
50	150	28.4%	34.9%	40.8%	58.1%	62.2%	65.3%	90.8%	93.4%	93.8%	99.8%	99.9%	99.9%
100	150	37.7%	44.8%	49.8%	66.5%	69.9%	72.8%	93.1%	95.1%	96.4%	100.0%	100.0%	100.0%
150	150	46.2%	51.1%	55.3%	67.1%	71.0%	74.3%	95.9%	97.0%	97.5%	100.0%	100.0%	100.0%
100	100	40.0%	46.1%	51.5%	65.4%	70.2%	73.3%	93.8%	96.3%	96.9%	100.0%	100.0%	100.0%
200	100	52.5%	57.3%	61.4%	71.6%	74.8%	77.0%	96.6%	97.3%	97.7%	100.0%	100.0%	100.0%
300	100	59.5%	63.7%	68.2%	75.0%	77.7%	80.0%	95.9%	97.1%	97.4%	100.0%	100.0%	100.0%
100	200	39.9%	46.9%	51.9%	66.2%	70.9%	73.2%	93.4%	94.8%	95.6%	99.8%	99.9%	99.9%
200	200	48.5%	54.8%	58.2%	68.4%	72.9%	76.2%	95.9%	97.0%	97.3%	100.0%	100.0%	100.0%
300	200	56.0%	59.9%	63.0%	69.3%	72.8%	75.9%	96.4%	97.4%	98.3%	100.0%	100.0%	100.0%
100	300	41.0%	47.4%	50.2%	67.4%	71.9%	73.4%	93.3%	94.9%	95.4%	100.0%	100.0%	100.0%
200	300	50.6%	55.6%	58.9%	68.7%	72.3%	74.4%	94.7%	95.8%	96.4%	100.0%	100.0%	100.0%
300	300	54.9%	59.0%	63.1%	72.3%	74.9%	77.3%	94.8%	96.8%	97.6%	100.0%	100.0%	100.0%
100	500	39.5%	45.0%	49.0%	65.1%	68.9%	71.2%	94.0%	95.6%	96.6%	99.9%	99.9%	99.9%
200	500	50.4%	54.4%	58.4%	69.4%	72.6%	75.6%	95.4%	97.2%	97.6%	100.0%	100.0%	100.0%
300	500	53.4%	58.3%	61.8%	71.2%	73.2%	75.2%	96.1%	97.4%	97.9%	100.0%	100.0%	100.0%

9 Extension

In this section, we relax Assumption B to allow for general weakly dependence idiosyncratic errors. Following Chamberlain and Rothschild (1983) we call a factor model with weak dependence idiosyncratic errors the approximate factor model. Approximate factor models are the primary research interests in a number of studies, e.g., Bai and Ng (2002), Bai (2003) and Bai and Li (2016), among others. To relax Assumption B, we introduce the following assumption to control the heteroskedasticity and weak correlations over cross section and time.

Assumption B'': (weak dependence on errors)

B''.1 $E(e_{it}) = 0$, and $E(e_{it}^8) \leq C$.

B''.2 Let $\mathbf{O}_t = E(e_t e_t')$, $\mathbf{O} = \frac{1}{T} \sum_{t=1}^T \mathbf{O}_t$, and $\mathbf{W} = \text{diag}(\mathbf{O})$, which is the diagonal matrix that sets the off-diagonal elements of \mathbf{O} to zero. Specifically, let w_i^2 be the i th diagonal element of \mathbf{W} , then $\mathbf{W} = \text{diag}(w_1^2, w_2^2, \dots, w_N^2)$.

B''.3 For all i , $C^{-2} \leq w_i^2 \leq C^2$;

B''.4 Let $\tau_{ij,t} \equiv E(e_{it} e_{jt})$, assume there exists some positive τ_{ij} such that $|\tau_{ij,t}| \leq \tau_{ij}$ for all t and $\sum_{i=1}^N \tau_{ij} \leq C$ for all j .

B''.5 Let $\rho_{i,ts} \equiv E(e_{it}e_{is})$, assume there exists some positive ρ_{ts} such that $|\rho_{i,ts}| \leq \rho_{ts}$ for all i and $\frac{1}{T} \sum_{t=1}^T \sum_{s=1}^T \rho_{ts} \leq C$.

B''.6 Assume $E \left[\left| \frac{1}{\sqrt{T}} \sum_{t=1}^T [e_{it}e_{jt} - E(e_{it}e_{jt})] \right|^4 \right] \leq C$ for all i and all j .

To be consistent with the changes in Assumption B'', we modify Assumptions C and D as follows.

Assumption C'':

C''.1 $\|\Lambda\| \leq C$ and $\|m_j\| \leq C$ for all j , where m_j is the transpose of the j th row of M .

C''.2 Let $\mathbb{P} = \Lambda' M' W^{-1} M \Lambda / N$, $\mathbb{R} = M' W^{-1} M / N$. We assume that $\mathbb{P}_\infty = \lim_{N \rightarrow \infty} \mathbb{P}$ and $\mathbb{R}_\infty = \lim_{N \rightarrow \infty} \mathbb{R}$ exist. Here \mathbb{P}_∞ and \mathbb{R}_∞ are some positive definite matrices.

Assumption D'': The estimator of w_j^2 for $j = 1, \dots, N$ takes value in a compact set: $[C^{-2}, C^2]$. Furthermore, M_{ff} is restricted to be in a set consisting of all semi-positive definite matrices with all elements bounded in the interval $[-C, C]$.

For theoretical analysis, we further assume the following two assumptions.

Assumption E'': We assume

E''.1 Let $\delta_{ijts} = E(e_{it}e_{js})$, and we assume $\frac{1}{NT} \sum_{i=1}^N \sum_{j=1}^N \sum_{t=1}^T \sum_{s=1}^T |\delta_{ijts}| \leq C$.

E''.2 Let $\pi_1 = \frac{1}{NT} \sum_{i=1}^N \sum_{j=1}^N \sum_{t=1}^T \sum_{s=1}^T \frac{\delta_{ijts}}{w_i^2 w_j^2} (m_i \otimes f_t)(m'_j \otimes f'_s)$, and assume $\lim_{N, T \rightarrow \infty} \pi_1 = \pi_{1\infty} > 0$; in other words, the limit of π_1 exists and is positive definite.

E''.3 Let $\pi_2 = \frac{1}{NT} \sum_{i=1}^N \sum_{j=1}^N \sum_{t=1}^T \sum_{s=1}^T \frac{q_{ijts}}{w_i^4 w_j^4} (m_i \otimes m_i)(m'_j \otimes m'_j)$ with $q_{ijts} = E[(e_{it}^2 - w_i^2)(e_{js}^2 - w_j^2)]$. We assume $\lim_{N, T \rightarrow \infty} \pi_2 = \pi_{2\infty} > 0$.

E''.4 Let $\pi_3 = \frac{1}{NT} \sum_{i=1}^N \sum_{j=1}^N \sum_{t=1}^T \sum_{s=1}^T \frac{\vartheta_{ijts}}{w_i^2 w_j^4} (m_i \otimes f_t)(m'_j \otimes m'_j)$ with $\vartheta_{ijts} = E[e_{it}(e_{js}^2 - w_j^2)]$. We assume $\lim_{N, T \rightarrow \infty} \pi_3 = \pi_{3\infty} > 0$.

E''.5 For each i , as $T \rightarrow \infty$, $\frac{1}{\sqrt{T}} \sum_{t=1}^T (e_{it}^2 - w_i^2) \xrightarrow{d} N(0, \omega_{i\infty}^2)$, with $\omega_{i\infty}^2 = \lim_{T \rightarrow \infty} \omega_i^2$ and $\omega_i^2 = \frac{1}{T} \sum_{t=1}^T \sum_{s=1}^T E[(e_{it}^2 - w_i^2)(e_{is}^2 - w_i^2)]$.

Assumption F'': We assume

F''.1 For all j , $E \left[\left\| \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^T \frac{m'_i \Lambda}{w_i^2} [e_{it}e_{jt} - E(e_{it}e_{jt})] \right\|^2 \right] \leq C$.

F''.2 We assume $E \left[\left\| \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^T \frac{m'_i \Lambda \Lambda' m_i}{w_i^2} (e_{it}^2 - w_i^2) \right\|^2 \right] \leq C$.

F''.3 For all t , $E \left[\left\| \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{s=1}^T \frac{1}{w_i^4} f_s [e_{it}e_{is} - E(e_{it}e_{is})] \right\|^2 \right] \leq C$.

F''.4 For all t , $E \left[\frac{1}{N} \sum_{i=1}^N \left\| \frac{1}{\sqrt{T}} \sum_{s=1}^T f_s [e_{it}e_{is} - E(e_{it}e_{is})] \right\|^2 \right] \leq C$.

F''.5 For all t , $E \left[\left\| \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{s=1}^T \frac{1}{w_i^4} m'_i \Lambda (e_{is}^2 - w_i^2) e_{it} \right\|^2 \right] \leq C$.

F'' .6 We assume $E \left[\left\| \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^T \sum_{s=1}^T \frac{1}{w_i^4} f_t e_{it} (e_{is}^2 - w_i^2) m'_i \right\|^2 \right] \leq C$.

Assumption E'' is used in deriving the limiting distributions. Assumption F'' provides some moment conditions which are needed in inferential analysis.

To remove the rotational indeterminacy, the identification conditions considered here, which are denoted by IC'', are the same with those in Section 3 except that the matrix Σ_{ee} is replaced with \mathbb{W} .

Even that the model allows for general weak dependence among idiosyncratic errors, we still use (3.2) as the objective function to estimate the loadings and idiosyncratic variances, with Σ_{ee} replaced by \mathbb{W} . Now the parameter is $\theta = (\Lambda, \mathbb{W})$. As shown in Bai and Li (2016), although the objective function is misspecified, the consistency of the estimated loadings can be maintained if some regularity conditions are satisfied.

Let $\hat{\theta} = (\hat{\Lambda}, \hat{\mathbb{W}})$ be the maximizer of the objective function. Then we can derive the first order conditions for Λ and \mathbb{W} , which are similar to (3.3) and (3.4), except that $\hat{\Sigma}_{ee}$ should be replaced by $\hat{\mathbb{W}}$. Based on these first order conditions, together with the similar arguments, we develop inferential theories under the weak dependence idiosyncratic errors. The following theorem presents the convergence rates of the MLE. The consistency is implied by the theorem.

Theorem 9.1 (Convergence rates) *Under Assumptions A, B'', C'', D'' and F'', together with IC'', when $N, T \rightarrow \infty$, we have*

$$\hat{\Lambda} - \Lambda = O_p \left(\frac{1}{\sqrt{NT}} \right) + O_p \left(\frac{1}{T} \right) + O_p \left(\frac{1}{N} \right), \quad \frac{1}{N} \sum_{i=1}^N (\hat{w}_i^2 - w_i^2)^2 = O_p \left(\frac{1}{T} \right) + O_p \left(\frac{1}{N^2} \right).$$

In contrast with the results in Theorem 4.1, we see that there is an extra term $O_p(\frac{1}{N})$ in $(\hat{\Lambda} - \Lambda)$ and another extra term $O_p(\frac{1}{N^2})$ in $\frac{1}{N} \sum_{i=1}^N (\hat{w}_i^2 - w_i^2)^2$ under the weak dependence data structure.

Before we state the asymptotic result of $\hat{\Lambda}$, below we first introduce some symbols.

$$\begin{aligned} \mathbb{D}_1^\dagger &= \left[\mathcal{D}[(\mathbb{P} \otimes I_r) + (I_r \otimes \mathbb{P})K_r] \right]', \\ \mathbb{B}_1^\dagger &= K_{kr}[(\mathbb{P}^{-1}\Lambda') \otimes \Lambda] + \mathbb{R}^{-1} \otimes I_r - K_{kr}(I_r \otimes \Lambda)(\mathbb{D}_1^\dagger)^{-1}\mathbb{D}_2[(\mathbb{P}^{-1}\Lambda') \otimes I_r], \\ \mathbb{B}_2^\dagger &= K_{kr}(I_r \otimes \Lambda)(\mathbb{D}_1^\dagger)^{-1}\mathbb{D}_3(\Lambda \otimes \Lambda)', \quad \mathbb{B}_3^\dagger = K_{kr}(I_r \otimes \Lambda)(\mathbb{D}_1^\dagger)^{-1}\mathbb{D}_3(\Lambda \otimes \Lambda)', \\ \mathbb{B}_4^\dagger &= \left((\mathbb{R}^{-1}) \otimes (\mathbb{P}^{-1}\Lambda') \right) - \frac{1}{2}K_{kr}(I_r \otimes \Lambda)(\mathbb{D}_1^\dagger)^{-1}\mathbb{D}_2(\mathbb{P} \otimes \mathbb{P})^{-1}(\Lambda \otimes \Lambda)', \\ \Delta^\dagger &= \mathbb{B}_2^\dagger \frac{1}{N} \sum_{i=1}^N \sum_{t=1}^T \frac{\varphi_i^2}{w_i^6} (m_i \otimes m_i), \\ \Pi^\dagger &= \mathbb{B}_4^\dagger \frac{1}{N} \sum_{i=1}^N \sum_{j=1, j \neq i}^N \frac{\mathbb{O}_{ij}}{w_i^2 w_j^2} (m_j \otimes m_i) - \mathbb{B}_3^\dagger \frac{1}{N} \sum_{i=1}^N \frac{\zeta_i}{w_i^4} (m_i \otimes m_i). \end{aligned}$$

where D_r^+ , \mathcal{D} , K_r , K_{kr} , \mathbb{D}_2 and \mathbb{D}_3 are defined the same as in Theorem 4.2; \mathbb{P} and \mathbb{R} are defined in Assumption C''; \mathbb{O}_{ij} is the (i, j) th entry of matrix \mathbb{O} ; $\zeta_i = \frac{1}{N} m_i' \Lambda \mathbb{P}^{-1} \Lambda' M' \mathbb{W}^{-1} (\mathbb{O} -$

$\mathbb{W})\mathbb{W}^{-1}M\Lambda\mathbb{P}^{-1}\Lambda'm_i - 2m_i'\Lambda\mathbb{G}_N\Lambda'M'\mathbb{W}^{-1}(\mathbb{O} - \mathbb{W})_i$ where $\mathbb{G}_N = N\mathbb{G}$ with $\mathbb{G} = (I_r + \Lambda'M'\mathbb{W}^{-1}M\Lambda)^{-1}$ and $(\mathbb{O} - \mathbb{W})_i$ is the i th column of $(\mathbb{O} - \mathbb{W})$; $\omega_i^2 = \frac{1}{T} \sum_{t=1}^T \sum_{s=1}^T E[(e_{it}^2 - w_i^2)(e_{is}^2 - w_i^2)]$ is defined in Assumption E''.5; both ζ_i and ω_i^2 are scalars. Then we provide the asymptotic representation of $\hat{\Lambda}$ in the following theorem.

Theorem 9.2 (Asymptotic representation for $\hat{\Lambda}$) *Under assumptions of Theorem 9.1,*

$$\begin{aligned} \text{vec}(\hat{\Lambda}' - \Lambda') &= \mathbb{B}_1^\dagger \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{w_i^2} (m_i \otimes f_t) e_{it} - \mathbb{B}_2^\dagger \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{w_i^4} (m_i \otimes m_i) (e_{it}^2 - w_i^2) \\ &\quad + \frac{1}{T} \Delta^\dagger + \frac{1}{N} \Pi^\dagger + O_p\left(\frac{1}{N\sqrt{T}}\right) + O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T^{3/2}}\right) + O_p\left(\frac{1}{N^2}\right), \end{aligned} \quad (9.1)$$

where the symbols $\mathbb{B}_1^\dagger, \mathbb{B}_2^\dagger, \Delta^\dagger$ and Π^\dagger are defined in the preceding paragraph.

Given the above theorem, we have the following corollary.

Corollary 9.1 (Limiting distribution for $\hat{\Lambda}$) *Under assumptions of Theorem 9.1 and Assumption E'', as $N, T \rightarrow \infty, N/T^2 \rightarrow 0$ and $T/N^3 \rightarrow 0$, we have*

$$\sqrt{NT} \left[\text{vec}(\hat{\Lambda}' - \Lambda') - \frac{1}{T} \Delta^\dagger - \frac{1}{N} \Pi^\dagger \right] \xrightarrow{d} N(0, \Xi),$$

where $\Xi = \lim_{N \rightarrow \infty} \Xi_{NT}$, and

$$\Xi_{NT} = \mathbb{B}_1^\dagger \pi_1 \mathbb{B}_1^{\dagger'} + \mathbb{B}_2^\dagger \pi_2 \mathbb{B}_2^{\dagger'} - \mathbb{B}_1^\dagger \pi_3 \mathbb{B}_2^{\dagger'} - \mathbb{B}_2^\dagger \pi_3' \mathbb{B}_1^{\dagger'}$$

where \mathbb{B}_1^\dagger and \mathbb{B}_2^\dagger are defined the same as in Theorem 9.2; the symbols π_1, π_2 and π_3 are defined in Assumption E''. Furthermore, by Assumption E''.2, E''.3 and E''.4, we have

$$\Xi = \mathbb{B}_1^\dagger \pi_{1\infty} \mathbb{B}_1^{\dagger'} + \mathbb{B}_2^\dagger \pi_{2\infty} \mathbb{B}_2^{\dagger'} - \mathbb{B}_1^\dagger \pi_{3\infty} \mathbb{B}_2^{\dagger'} - \mathbb{B}_2^\dagger \pi_{3\infty}' \mathbb{B}_1^{\dagger'}$$

where the symbols $\pi_{1\infty}, \pi_{2\infty}$ and $\pi_{3\infty}$ are defined in Assumption E''.

we also have the following theorem for w_i^2 .

Theorem 9.3 (Asymptotic properties for \hat{w}_i^2) *Under assumptions of Theorem 9.1,*

$$\hat{w}_i^2 - w_i^2 = \frac{1}{T} \sum_{t=1}^T (e_{it}^2 - w_i^2) + O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T}\right) + O_p\left(\frac{1}{N}\right).$$

As $N, T \rightarrow \infty$ and $T/N^2 \rightarrow 0$, we have

$$\sqrt{T}(\hat{w}_i^2 - w_i^2) = \frac{1}{\sqrt{T}} \sum_{t=1}^T (e_{it}^2 - w_i^2) + o_p(1).$$

Furthermore, by Assumption E''.5, we have

$$\sqrt{T}(\hat{w}_i^2 - w_i^2) \xrightarrow{d} N(0, \omega_{i\infty}^2),$$

where $\omega_{i\infty}^2$ is defined in Assumption E''.5.

This limiting result is the same as that in the unconstrained approximate factor model, see Bai and Li (2016).

10 Conclusion

This paper considers the ML estimation of large dimensional constrained factor models in which both cross sectional units (N) and time periods (T) are large but the number of loadings is fixed. We investigate the asymptotic properties of the MLE including consistency, convergence rates, asymptotic representations and limiting distributions. We show that the MLE for the loadings in a constrained factor model converges much faster than that in a standard factor model. In addition, we also find that the MLE has a non-negligible bias asymptotically and some bias corrections are needed when conducting inference. A W statistic is proposed to conduct model specification check in a constrained factor model versus a standard factor model. The test is valid for a large N and a large T setup. We also analyze partially constrained factor models where only partial factor loadings are constrained. We run simulations to investigate the finite sample performance of the MLE and the proposed W test. The simulation results are encouraging and show that the MLE outperform the PC estimates and the proposed W test has good empirical sizes and powers. Monte carlo simulations show that our proposed MLE has better finite sample performances than that of PC estimates. In addition, we consider the extension of a general weak dependence structure on idiosyncratic errors and we study MLE asymptotic properties of the resulting approximate factor models.

Appendices: Proofs of the main theoretical results

We will prove the main theoretical results reported in Section 4 in appendices A and B. The supplementary appendices C to G contain proofs of additional results reported in the paper and also report some additional simulation results.

Appendix A: Proof for Proposition 4.1 (consistency)

The following notation will be used in this appendix.

$$\begin{aligned}\hat{P} &= \frac{1}{N} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda}; & \hat{R} &= \frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M; & \hat{G} &= (I_r + \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1}; \\ \hat{P}_N &= N \cdot \hat{P} = \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda}; & \hat{R}_N &= N \cdot \hat{R} = M' \hat{\Sigma}_{ee}^{-1} M, & \hat{G}_N &= N \cdot \hat{G}.\end{aligned}$$

From $(A + B)^{-1} = A^{-1} - A^{-1}B(A + B)^{-1}$, we have $\hat{P}_N^{-1} = \hat{G}(I - \hat{G})^{-1}$. From $\Sigma_{zz} = M\Lambda\Lambda'M' + \Sigma_{ee}$, we have

$$\Sigma_{zz}^{-1} = \Sigma_{ee}^{-1} - \Sigma_{ee}^{-1} M \Lambda (I_r + \Lambda' M' \Sigma_{ee}^{-1} M \Lambda)^{-1} \Lambda' M' \Sigma_{ee}^{-1}. \quad (\text{A.1})$$

It follows that

$$\hat{\Lambda}' M' \hat{\Sigma}_{zz}^{-1} = \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} - \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} (I_r + \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} = \hat{G} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1}. \quad (\text{A.2})$$

We use symbols with superscript “*” to denote the true parameters. Variables without superscript “*” denote the arguments of the likelihood function.

Let $\theta = (\Lambda, \sigma_1^2, \dots, \sigma_N^2)$ and let Θ be the parameter set such that Λ take values in a compact set and $C^{-2} \leq \sigma_i^2 \leq C^2$ for all $i = 1, \dots, N$. We assume $\theta^* = (\Lambda^*, \sigma_1^{*2}, \dots, \sigma_N^{*2})$ is an interior point of Θ . For simplicity, we write $\theta = (\Lambda, \Sigma_{ee})$ and $\theta^* = (\Lambda^*, \Sigma_{ee}^*)$.

The following lemmas are useful for our analysis

Lemma A.1 *Under assumptions of A-D, we have*

$$(a) \sup_{\theta \in \Theta} \frac{1}{NT} \left| \text{tr} \left[\Lambda^{*'} M' \Sigma_{zz}^{-1} \sum_{t=1}^T e_t f_t^{*'} \right] \right| \xrightarrow{p} 0;$$

$$(b) \sup_{\theta \in \Theta} \frac{1}{NT} \left| \text{tr} \left[\sum_{t=1}^T (e_t e_t' - \Sigma_{ee}^*) \Sigma_{zz}^{-1} \right] \right| \xrightarrow{p} 0;$$

where $\theta^* = (\Lambda^*, \Sigma_{ee}^*)$ denotes the true parameters and $\Sigma_{zz} = M \Lambda \Lambda' M' + \Sigma_{ee}$.

PROOF OF LEMMA A.1. First, we consider (a). Let m_{ip} be the (i, p) th element of M for $i = 1, \dots, N, p = 1, \dots, k$ and $\Lambda = [\lambda_1, \lambda_2, \dots, \lambda_k]'$. By equation (A.1), we have

$$\begin{aligned} \frac{1}{NT} \Lambda^{*'} M' \Sigma_{zz}^{-1} \sum_{t=1}^T e_t f_t^{*'} &= \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \left(\sum_{p=1}^k \lambda_p^* m_{ip} \right) \frac{1}{\sigma_i^2} e_{it} f_t^{*'} \\ &- \Lambda^{*'} M' \Sigma_{ee}^{-1} M \Lambda (I_r + \Lambda' M' \Sigma_{ee}^{-1} M \Lambda)^{-1} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \left(\sum_{p=1}^k \lambda_p m_{ip} \right) \frac{1}{\sigma_i^2} e_{it} f_t^{*'} \end{aligned} \quad (\text{A.3})$$

By the Cauchy-Schwartz inequality, the first term on the right side of (A.3) is bounded in norm by

$$\left(\frac{1}{N} \sum_{i=1}^N \frac{1}{\sigma_i^4} \left\| \sum_{p=1}^k \lambda_p^* m_{pi} \right\|^2 \right)^{1/2} \left[\frac{1}{N} \sum_{i=1}^N \left\| \frac{1}{T} \sum_{t=1}^T f_t^* e_{it} \right\|^2 \right]^{1/2}.$$

The first factor $\left(\frac{1}{N} \sum_{i=1}^N \frac{1}{\sigma_i^4} \left\| \sum_{p=1}^k \lambda_p^* m_{pi} \right\|^2 \right)^{1/2}$ is bounded by the boundedness of σ^{-2} and $\frac{1}{N} \sum_{i=1}^N \left\| \sum_{p=1}^k \lambda_p^* m_{pi} \right\|^2$ by Assumptions C and D. The second factor does not depend on any unknown parameters, and it is $O_p(T^{-1/2})$ because $E \left(\frac{1}{N} \sum_{i=1}^N \left\| \frac{1}{T} \sum_{t=1}^T f_t^* e_{it} \right\|^2 \right) = O(T^{-1})$. Therefore, the first part on the right hand side of (A.3) is $o_p(1)$ uniformly on θ . For the second part, we rewrite it in terms of P_N as

$$\Lambda^{*'} M' \Sigma_{ee}^{-1} M \Lambda P_N^{-1/2} (P_N^{-1} + I_r)^{-1} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T P_N^{-1/2} \left(\sum_{p=1}^k \lambda_p m_{ip} \right) \frac{1}{\sigma_i^2} e_{it} f_t^{*'} \quad (\text{A.4})$$

The term $\Lambda^{*'} M' \Sigma_{ee}^{-1} M \Lambda P_N^{-1/2} = \sum_{i=1}^N \frac{1}{\sigma_i^2} \left(\sum_{p=1}^k \lambda_p^* m_{ip} \right) \left(\sum_{p=1}^k \lambda_p' m_{ip} \right) P_N^{-1/2}$ is bounded in norm by

$$C \left(\sum_{i=1}^N \left\| \sum_{p=1}^k \lambda_p^* m_{ip} \right\|^2 \right)^{1/2} \left(\sum_{i=1}^N \frac{1}{\sigma_i^2} \left\| \sum_{p=1}^k \lambda_p' m_{ip} P_N^{-1/2} \right\|^2 \right)^{1/2} = a_1, \quad \text{say.}$$

Notice that

$$\begin{aligned} \sum_{i=1}^N \frac{1}{\sigma_i^2} \left\| P_N^{-1/2} \sum_{p=1}^k \lambda_p m_{ip} \right\|^2 &= \sum_{i=1}^N \frac{1}{\sigma_i^2} \left(\sum_{p=1}^k \lambda_p' m_{ip} P_N^{-1} \sum_{q=1}^k \lambda_q m_{iq} \right) \\ &= \text{tr} \left[P_N^{-1} \Lambda' M' \Sigma_{ee}^{-1} M \Lambda \right] = \text{tr} [P_N^{-1} P_N] = r. \end{aligned} \quad (\text{A.5})$$

We have $a_1 = O_p(N^{1/2})$. As regard to the term $\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T P_N^{-1/2} (\sum_{p=1}^k \lambda_p m_{ip}) \frac{1}{\sigma_i^2} e_{it} f_t^{*}$, it is bounded in norm by

$$C \frac{1}{\sqrt{N}} \left(\sum_{i=1}^N \frac{1}{\sigma_i^2} \left\| P_N^{-1/2} \sum_{p=1}^k \lambda_p m_{ip} \right\|^2 \right)^{1/2} \left(\frac{1}{N} \sum_{i=1}^N \left\| \frac{1}{T} \sum_{t=1}^T f_t^* e_{it} \right\|^2 \right)^{1/2} = O_p(N^{-1/2} T^{-1/2})$$

by (A.5). In addition, the term $(P_N^{-1} + I_r)^{-1} = O_p(1)$ uniformly on Θ . So the expression in (A.4) is $O_p(T^{-1/2})$ uniformly on θ . Then result (a) follows.

Next, we consider (b). By equation (A.1), we have

$$\begin{aligned} & \text{tr} \left[\frac{1}{NT} \sum_{t=1}^T (e_t e_t' - \Sigma_{ee}^*) \Sigma_{zz}^{-1} \right] \\ &= \text{tr} \left[\frac{1}{NT} \sum_{t=1}^T (e_t e_t' - \Sigma_{ee}^*) (\Sigma_{ee}^{-1} - \Sigma_{ee}^{-1} M \Lambda (I_r + \Lambda' M' \Sigma_{ee}^{-1} M \Lambda)^{-1} \Lambda' M' \Sigma_{ee}^{-1}) \right] \\ &= \text{tr} \left[\frac{1}{NT} \sum_{t=1}^T (e_t e_t' - \Sigma_{ee}^*) \Sigma_{ee}^{-1} \right] \\ &\quad - \text{tr} \left[\frac{1}{NT} \sum_{t=1}^T (\Lambda' M' \Sigma_{ee}^{-1} (e_t e_t' - \Sigma_{ee}^*) \Sigma_{ee}^{-1} M \Lambda) (I_r + \Lambda' M' \Sigma_{ee}^{-1} M \Lambda)^{-1} \right]. \end{aligned}$$

The first term $\text{tr} \left[\frac{1}{NT} \sum_{t=1}^T (e_t e_t' - \Sigma_{ee}^*) \Sigma_{ee}^{-1} \right] = \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} (e_{it}^2 - \sigma_i^{*2})$ is bounded by

$$\left(\frac{1}{N} \sum_{i=1}^N \frac{1}{\sigma_i^4} \right)^{1/2} \left(\frac{1}{N} \sum_{i=1}^N \left(\frac{1}{T} \sum_{t=1}^T e_{it}^2 - \sigma_i^{*2} \right)^2 \right)^{1/2},$$

which is $O_p(T^{-1/2})$ uniformly on θ . The second term can be written as

$$\text{tr} \left[\frac{1}{NT} P_N^{-1/2} \Lambda' M' \Sigma_{ee}^{-1} \left[\sum_{t=1}^T (e_t e_t' - \Sigma_{ee}^*) \right] \Sigma_{ee}^{-1} M \Lambda P_N^{-1/2} (P_N^{-1} + I_r)^{-1} \right].$$

The above term is equal to

$$\text{tr} \left[\left(\frac{1}{NT} \sum_{i=1}^N \sum_{j=1}^N \frac{1}{\sigma_i^2 \sigma_j^2} P_N^{-1/2} \sum_{p=1}^k \lambda_p m_{ip} \sum_{q=1}^k \lambda'_q m_{qj} P_N^{-1/2} \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})] \right) (P_N^{-1} + I_r)^{-1} \right].$$

Since the expression

$$\frac{1}{NT} \sum_{i=1}^N \sum_{j=1}^N \frac{1}{\sigma_i^2 \sigma_j^2} P_N^{-1/2} \sum_{p=1}^k \lambda_p m_{ip} \sum_{q=1}^k \lambda'_q m_{qj} P_N^{-1/2} \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})]$$

is bounded in norm by

$$C^2 \left[\sum_{i=1}^N \frac{1}{\sigma_i^2} \left\| P_N^{-1/2} \sum_{p=1}^k \lambda_p m_{ip} \right\|^2 \right] \left[\frac{1}{N^2} \sum_{i=1}^N \sum_{j=1}^N \left(\frac{1}{T} \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})] \right)^2 \right]^{1/2}$$

which is $O_p(T^{-1/2})$ uniformly on θ by (A.5). Given $(P_N^{-1} + I_r)^{-1} = O(1)$ uniformly on θ , the second term is $o_p(1)$ uniformly on θ . This proves (b). \square

Lemma A.2 *Under Assumptions A-D, we have*

$$(a) \quad \left\| \frac{1}{N} \Lambda^{*'} M' (\hat{\Sigma}_{ee}^{-1} - \Sigma_{ee}^{*-1}) M \Lambda^* \right\| = O_p \left(\left[\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^{*2})^2 \right]^{\frac{1}{2}} \right);$$

$$(b) \quad \left\| \frac{1}{N} M' (\hat{\Sigma}_{ee}^{-1} - \Sigma_{ee}^{*-1}) M \right\| = O_p \left(\left[\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^{*2})^2 \right]^{\frac{1}{2}} \right).$$

Given the above results, if $N^{-1} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^{*2})^2 = o_p(1)$, we have

$$(c) \quad \hat{R}_N = O_p(N), \quad \hat{R} = \frac{1}{N} \hat{R}_N = O_p(1);$$

$$(d) \quad \|\hat{R}^{-1/2}\| = O_p(1).$$

where \hat{R} and \hat{R}_N are defined above appendix A.

PROOF OF LEMMA A.2. We first consider (a). The left hand side of (a) can be written as

$$\frac{1}{N} \sum_{i=1}^N \left(\sum_{p=1}^k \lambda_p^* m_{ip} \right) \left(\sum_{q=1}^k m_{qi} \lambda_q^{*'} \right) \frac{\hat{\sigma}_i^2 - \sigma_i^{*2}}{\hat{\sigma}_i^2 \sigma_i^{*2}},$$

which is bounded in norm by

$$C^4 \left(\frac{1}{N} \sum_{i=1}^N \left\| \sum_{p=1}^k \lambda_p^* m_{ip} \right\|^4 \right)^{1/2} \left(\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^{*2})^2 \right)^{1/2}.$$

Then result (a) follows because $\left\| \sum_{p=1}^k \lambda_p^* m_{ip} \right\|^4$ is bounded by Assumption C.

Next, we consider (b). The left hand side of (b) can be written as $\frac{1}{N} \sum_{i=1}^N m_i m_i' \frac{\hat{\sigma}_i^2 - \sigma_i^{*2}}{\hat{\sigma}_i^2 \sigma_i^{*2}}$, where m_i is the transpose of the i th row of M . This term is bounded in norm by

$$C^4 \left(\frac{1}{N} \sum_{i=1}^N \|m_i\|^4 \right)^{1/2} \left(\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^{*2})^2 \right)^{1/2}.$$

Then result (b) follows because $\frac{1}{N} \sum_{i=1}^N \|m_i\|^4$ is bounded by Assumption C.

We now consider (c). From result (b) and result $N^{-1} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^{*2})^2 = o_p(1)$, we have $\hat{R} - \frac{1}{N} M' \Sigma_{ee}^{-1} M = o_p(1)$ which implies $\hat{R} \xrightarrow{P} R > 0$, where R is defined in Assumption C. So $\hat{R} = O_p(1)$ and $\hat{R}_N = N\hat{R} = O_p(N)$. Result (c) follows.

Result (d) is a direct result of $\|\hat{R}^{-1/2}\|^2 = \text{tr}(\hat{R}^{-1}) = O_p(1)$ by $\hat{R} \xrightarrow{P} R > 0$ from result (c).

This completes the proof of Lemma A.2. \square

Lemma A.3 *Under Assumptions A-D, we have*

$$(a) \quad \frac{1}{N^2} \hat{P}^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T (e_t e_t' - \Sigma_{ee}) \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}^{-1} = \|\hat{P}^{-1/2}\|^2 \cdot O_p(T^{-1/2});$$

$$(b) \quad \frac{1}{N} \hat{P}^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T e_t f_t' = \|\hat{P}^{-1/2}\| \cdot O_p(T^{-1/2});$$

- (c) $\frac{1}{N^2} \hat{P}^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} (\hat{\Sigma}_{ee} - \Sigma_{ee}) \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}^{-1} = \|\hat{P}_N^{-1}\| \cdot O_p(1);$
- (d) $\frac{1}{NT} \sum_{t=1}^T f_t e_t' \hat{\Sigma}_{ee}^{-1} M \hat{R}^{-1} = O_p(T^{-1/2});$
- (e) $\frac{1}{N^2} \hat{P}^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T [e_t e_t' - \Sigma_{ee}] \hat{\Sigma}_{ee}^{-1} M \hat{R}^{-1} = \|\hat{P}^{-1/2}\| \cdot O_p(T^{-1/2});$
- (f) $\frac{1}{N^2} \hat{P}^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} (\hat{\Sigma}_{ee} - \Sigma_{ee}) \hat{\Sigma}_{ee}^{-1} M \hat{R}^{-1} = \|\hat{P}^{-1/2}\| \cdot O_p\left(\left[\frac{1}{N^3} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2\right]^{\frac{1}{2}}\right).$

PROOF OF LEMMA A.3. We first consider (a). The left hand side can be rewritten as

$$\frac{1}{N^2} \hat{P}^{-1/2} \left[\sum_{i=1}^N \sum_{j=1}^N \hat{P}^{-1/2} \left(\sum_{p=1}^k \hat{\lambda}_p m_{ip} \right) \frac{1}{\hat{\sigma}_i^2 \hat{\sigma}_j^2} \frac{1}{T} \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})] \left(\sum_{q=1}^k m_{jq} \hat{\lambda}'_q \right) \hat{P}^{-1/2} \right] \hat{P}^{-1/2},$$

which is bounded in norm by

$$C^2 \|\hat{P}^{-1/2}\|^2 \left[\sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2} \left\| \hat{P}_N^{-1/2} \sum_{p=1}^k \hat{\lambda}_p m_{ip} \right\|^2 \right] \left[\frac{1}{N^2} \sum_{i=1}^N \sum_{j=1}^N \left| \frac{1}{T} \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})] \right|^2 \right]^{1/2},$$

which is $\|\hat{P}^{-1/2}\|^2 \cdot O_p(T^{-1/2})$ by (A.5). Thus, (a) follows.

Next, we consider (b). The left hand side can be rewritten as

$$\frac{1}{\sqrt{N}} \hat{P}^{-1/2} \sum_{i=1}^N \hat{P}_N^{-1/2} \frac{1}{\hat{\sigma}_i^2} \sum_{p=1}^k \hat{\lambda}_p m_{ip} \frac{1}{T} \sum_{t=1}^T e_{it} f'_t,$$

which is bounded in norm by

$$C \|\hat{P}^{-1/2}\| \left(\sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2} \left\| \hat{P}_N^{-1/2} \sum_{p=1}^k \hat{\lambda}_p m_{ip} \right\|^2 \right)^{1/2} \left(\frac{1}{N} \sum_{i=1}^N \left\| \frac{1}{T} \sum_{t=1}^T e_{it} f'_t \right\|^2 \right)^{1/2},$$

which is $\|\hat{P}^{-1/2}\| \cdot O_p(T^{-1/2})$ by (A.5). This proves result (b).

To prove result (c), notice that $\hat{\Sigma}_{ee}^{-1} (\hat{\Sigma}_{ee} - \Sigma_{ee})$ is bounded by $2C^4 I_N$ by $C^{-2} \leq \hat{\sigma}_i^2 \leq C^2$ and $C^{-2} \leq \sigma_i^2 \leq C^2$. Hence, the left hand side is bounded in norm by

$$\left\| \hat{P}_N^{-1} \hat{\Lambda}' M' (2C^4 I_N) \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}_N^{-1} \right\| = 2C^4 \|\hat{P}_N^{-1}\|.$$

Result (c) then follows.

We now consider (d). The left hand side is equal to

$$\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\hat{\sigma}_i^2} f_t e_{it} m'_i \hat{R},$$

which is bounded in norm by

$$C \|\hat{R}\| \cdot \left[\frac{1}{N} \sum_{i=1}^N \|m_i\|^2 \right]^{1/2} \left[\frac{1}{N} \sum_{i=1}^N \left\| \frac{1}{T} \sum_{t=1}^T f_t e_{it} \right\|^2 \right]^{1/2},$$

which is $O_p(T^{-1/2})$ by Lemma A.2(c) and Assumption C. Hence, result (d) follows.

For result (e), the left hand side is equal to

$$\frac{1}{N^{3/2}} \hat{P}^{-1/2} \left[\sum_{i=1}^N \sum_{j=1}^N \hat{P}_N^{-1/2} \left(\sum_{p=1}^k \hat{\lambda}_p m_{ip} \right) \frac{1}{\hat{\sigma}_i^2 \hat{\sigma}_j^2} \frac{1}{T} \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})] m'_j \right] \hat{R}^{-1},$$

which is bounded in norm by

$$\begin{aligned} & C^2 \|\hat{P}^{-1/2}\| \cdot \|\hat{R}^{-1}\| \cdot \left[\sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2} \left\| \hat{P}_N^{-1/2} \sum_{p=1}^k \hat{\lambda}_p m_{ip} \right\|^2 \right]^{1/2} \left[\frac{1}{N} \sum_{j=1}^N \|m_j\|^2 \right]^{1/2} \\ & \times \left[\frac{1}{N^2} \sum_{i=1}^N \sum_{j=1}^N \left| \frac{1}{T} \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})] \right|^2 \right]^{1/2}, \end{aligned}$$

which is $\|\hat{P}^{-1/2}\| \cdot O_p(T^{-1/2})$ by (A.5) and Lemma A.2(c). Thus, result (d) follows.

Finally, we consider (f). The left hand side can be written as

$$\frac{1}{N^{3/2}} \hat{P}^{-1/2} \sum_{i=1}^N \hat{P}_N^{-1/2} \left(\sum_{p=1}^k \hat{\lambda}_p m_{ip} \right) \left(\frac{\hat{\sigma}_i^2 - \sigma_i^2}{\hat{\sigma}_i^4} \right) m'_i \hat{R}^{-1},$$

which is bounded in norm by

$$\frac{1}{N} \cdot \|\hat{P}^{-1/2}\| \cdot \|\hat{R}^{-1}\| \left[\sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2} \left\| \hat{P}_N^{-1/2} \sum_{p=1}^k \hat{\lambda}_p m_{ip} \right\|^2 \right]^{1/2} \left[\frac{1}{N} \sum_{i=1}^N \frac{\|m_i\|^2}{\hat{\sigma}_i^4} (\hat{\sigma}_i^2 - \sigma_i^2)^2 \right]^{1/2}.$$

By the boundedness of $\|m_i\|$ and $\hat{\sigma}^{-2}$ by Assumptions C and D, we have

$$\frac{1}{N} \sum_{i=1}^N \frac{\|m_i\|^2}{\hat{\sigma}_i^4} (\hat{\sigma}_i^2 - \sigma_i^2)^2 \leq C \frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2.$$

This result, together with (A.5) and Lemma A.2(c), gives result (f). \square

PROOF OF PROPOSITION 4.1. Throughout the proof, we use the following centered objective function

$$L(\theta) = \bar{L}(\theta) + R(\theta),$$

where

$$\bar{L}(\theta) = -\frac{1}{N} \ln |\Sigma_{zz}| - \frac{1}{N} \text{tr} \left(\Sigma_{zz}^* \Sigma_{zz}^{-1} \right) + 1 + \frac{1}{N} \ln |\Sigma_{zz}^*|$$

and

$$R(\theta) = -\frac{1}{N} \text{tr} \left[(M_{zz} - \Sigma_{zz}^*) \Sigma_{zz}^{-1} \right],$$

where $\Sigma_{zz} = M \Lambda \Lambda' M' + \Sigma_{ee}$ and $\Sigma_{zz}^* = M \Lambda^* \Lambda^{*'} M' + \Sigma_{ee}^*$. The above objective function differs from the objective function of the main text only by a constant and is convenient for the subsequent analysis. By the definition of M_{zz} , we have

$$R(\theta) = -2 \frac{1}{NT} \text{tr} \left[M \Lambda^* \sum_{t=1}^T f_t^* e_t' \Sigma_{zz}^{-1} \right] - \frac{1}{NT} \text{tr} \left[\sum_{t=1}^T (e_t e_t' - \Sigma_{ee}^*) \Sigma_{zz}^{-1} \right].$$

By Lemma A.1, we have $\sup_{\theta} |R(\theta)| = o_p(1)$. Since $\hat{\theta}$ maximizes $L(\theta)$, it follows $\bar{L}(\hat{\theta}) + R(\hat{\theta}) \geq \bar{L}(\theta^*) + R(\theta^*)$. This implies that $\bar{L}(\hat{\theta}) \geq \bar{L}(\theta^*) + R(\theta^*) - R(\hat{\theta}) \geq \bar{L}(\theta^*) - 2 \sup_{\theta \in \Theta} |R(\theta)| = -|o_p(1)|$, where $\bar{L}(\theta^*)$ is normalized to be zero.

Now consider $\bar{L}(\hat{\theta})$ which is equivalent to

$$\bar{L}(\hat{\theta}) = -\frac{1}{N} \ln |\hat{\Sigma}_{zz}| - \frac{1}{N} \text{tr}(\Sigma_{zz}^* \hat{\Sigma}_{zz}^{-1}) + 1 + \frac{1}{N} \ln |\Sigma_{zz}^*|. \quad (\text{A.6})$$

By $\Sigma_{zz} = M\Lambda\Lambda'M' + \Sigma_{ee}$, we have $|\Sigma_{zz}| = |\Sigma_{ee}| \cdot |I_r + \Lambda'M'\Sigma_{ee}^{-1}M\Lambda|$. Similarly, $|\Sigma_{zz}^*| = |\Sigma_{ee}^*| \cdot |I_r + \Lambda^*M'\Sigma_{ee}^{*-1}M\Lambda^*|$. Then equation (A.6) can be written as

$$\begin{aligned} \bar{L}(\hat{\theta}) &= -\frac{1}{N} \ln |\hat{\Sigma}_{ee}| - \frac{1}{N} \ln |I_r + \hat{\Lambda}'M'\Sigma_{ee}^{-1}M\hat{\Lambda}| - \frac{1}{N} \text{tr}[M\Lambda^* \Lambda^*M'\hat{\Sigma}_{zz}^{-1}] \\ &\quad - \frac{1}{N} \text{tr}[\Sigma_{ee}^* \hat{\Sigma}_{zz}^{-1}] + \frac{1}{N} \ln |\Sigma_{ee}^*| + \frac{1}{N} \ln |I_r + \Lambda^*M'\Sigma_{ee}^{*-1}M\Lambda^*| + 1 \\ &= \left\{ -\frac{1}{N} \ln |\hat{\Sigma}_{ee}| + \frac{1}{N} \ln |\Sigma_{ee}^*| - \frac{1}{N} \text{tr}[\Sigma_{ee}^* \hat{\Sigma}_{zz}^{-1}] + 1 \right\} \\ &\quad + \left\{ -\frac{1}{N} \text{tr}[M\Lambda^* \Lambda^*M'\hat{\Sigma}_{zz}^{-1}] \right\} + \left\{ -\frac{1}{N} \ln |I_r + \hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}| \right\} \\ &\quad + \left\{ \frac{1}{N} \ln |I_r + \Lambda^*M'\Sigma_{ee}^{*-1}M\Lambda^*| \right\}. \end{aligned}$$

Notice that

$$\frac{1}{N} \text{tr}[\Sigma_{ee}^* \hat{\Sigma}_{zz}^{-1}] = \frac{1}{N} \text{tr}[\Sigma_{ee}^* \hat{\Sigma}_{ee}^{-1}] - \frac{1}{N} \text{tr}[\Sigma_{ee}^* \hat{\Sigma}_{ee}^{-1} M\hat{\Lambda}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}] = \frac{1}{N} \text{tr}[\Sigma_{ee}^* \hat{\Sigma}_{ee}^{-1}] + o_p(1)$$

by

$$0 < \frac{1}{N} \text{tr}[\Sigma_{ee}^* \hat{\Sigma}_{ee}^{-1} M\hat{\Lambda}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}] \leq C \frac{1}{N} \text{tr}[\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}\hat{\Lambda}] \leq C \frac{r}{N},$$

where we use the fact that there exists a constant C such that $\Sigma_{ee}^* \hat{\Sigma}_{ee}^{-1} \leq C \cdot I_N$ due to the boundedness of $\hat{\sigma}_i^2$ and σ_i^{*2} .

Given the above result, together with $\frac{1}{N} \ln |I_r + \Lambda^*M'\Sigma_{ee}^{*-1}M\Lambda^*| = O(\ln N/N)$, we can further write $\bar{L}(\hat{\theta})$ as

$$\begin{aligned} \bar{L}(\hat{\theta}) &= - \left\{ \frac{1}{N} \ln |\hat{\Sigma}_{ee}| - \frac{1}{N} \ln |\Sigma_{ee}^*| + \frac{1}{N} \text{tr}[\Sigma_{ee}^* \hat{\Sigma}_{ee}^{-1}] - 1 \right\} \\ &\quad - \left\{ \frac{1}{N} \text{tr}[M\Lambda^* \Lambda^*M'\hat{\Sigma}_{zz}^{-1}] \right\} - \left\{ \frac{1}{N} \ln |I_r + \hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}| \right\} + o_p(1). \end{aligned}$$

The above three expressions in the big curly bracket are all non-negative. Together with $\bar{L}(\hat{\theta}) \geq -2|o_p(1)|$, we have that each expression is $o_p(1)$, that is,

$$\frac{1}{N} \ln |\hat{\Sigma}_{ee}| - \frac{1}{N} \ln |\Sigma_{ee}^*| + \frac{1}{N} \text{tr}[\Sigma_{ee}^* \hat{\Sigma}_{ee}^{-1}] - 1 \xrightarrow{p} 0, \quad (\text{A.7})$$

$$\frac{1}{N} \text{tr}[M\Lambda^* \Lambda^*M'\hat{\Sigma}_{zz}^{-1}] \xrightarrow{p} 0. \quad (\text{A.8})$$

Equation (A.7) is equivalent to

$$\frac{1}{N} \sum_{i=1}^N (\ln \hat{\sigma}_i^2 - \ln \sigma_i^{*2} + \frac{\sigma_i^{*2}}{\hat{\sigma}_i^2} - 1) \xrightarrow{p} 0.$$

Consider the function $g(x) = \ln x + \frac{\sigma_i^{*2}}{x} - \ln \sigma_i^{*2} - 1$. Given that $0 < C^{-2} \leq \sigma_i^2 \leq C^2 < \infty$ for $C > 1$, for any $x \in [C^{-2}, C^2]$, we can find a constant d (for example, let $d = \frac{1}{4C^4}$) such that $g(x) \geq d(x - \sigma_i^{*2})^2$. It follows

$$o_p(1) = \frac{1}{N} \sum_{i=1}^N (\ln \hat{\sigma}_i^2 + \frac{\sigma_i^{*2}}{\hat{\sigma}_i^2} - 1 - \ln \sigma_i^{*2}) \geq d \frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^{*2})^2.$$

The above argument implies

$$\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^{*2})^2 \xrightarrow{p} 0. \quad (\text{A.9})$$

This proves the first result of Proposition 4.1.

Next, we consider (A.8), which is equivalent to

$$\frac{1}{N} \text{tr}(M\Lambda^* \Lambda'^* M' \hat{\Sigma}_{zz}^{-1}) = \frac{1}{N} \text{tr} \left[\Lambda'^* M' (\hat{\Sigma}_{ee}^{-1} - \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{G} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1}) M \Lambda^* \right].$$

By $(I_r + \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} = (\hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} - (\hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} (I_r + \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1}$, the preceding expression can be alternatively written as

$$\begin{aligned} & \frac{1}{N} \text{tr}(M\Lambda^* \Lambda'^* M' \hat{\Sigma}_{zz}^{-1}) \\ &= \frac{1}{N} \text{tr} \left[\Lambda'^* M' \hat{\Sigma}_{ee}^{-1} M \Lambda^* - \Lambda'^* M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} (\hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \Lambda^* \right] \\ & \quad + \frac{1}{N} \text{tr} \left[\Lambda'^* M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} (\hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} (I_r + \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \Lambda^* \right] \end{aligned}$$

Both terms on the right hand side are non-negative. By (A.8), it follows that

$$\frac{1}{N} \text{tr} \left[\Lambda'^* M' \hat{\Sigma}_{ee}^{-1} M \Lambda^* - \Lambda'^* M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} (\hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \Lambda^* \right] \xrightarrow{p} 0, \quad (\text{A.10})$$

$$\frac{1}{N} \text{tr} \left[\Lambda'^* M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} (\hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} (I_r + \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \Lambda^* \right] \xrightarrow{p} 0. \quad (\text{A.11})$$

By (A.9) and Lemma A.2(a), we know $\frac{1}{N} \text{tr}(\Lambda'^* M' \hat{\Sigma}_{ee}^{-1} M \Lambda^*)$ converges to a positive constant. Then (A.10) implies that $\frac{1}{N} \text{tr}(\Lambda'^* M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} (\hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \Lambda^*)$ converges to the same positive constant. Together with (A.11), we have $(I_r + \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} = o_p(1)$, i.e. $\hat{G} = o_p(1)$. Furthermore, from $\hat{P}_N^{-1} = \hat{G}(I - \hat{G})^{-1}$, we have $\hat{P}_N^{-1} = o_p(1)$. We obtain the following results

$$\hat{G} = o_p(1); \quad \hat{P}_N^{-1} = o_p(1). \quad (\text{A.12})$$

Consider (A.10) again. The matrix on the left-hand side is finite dimensional ($r \times r$) and is semi-positive definite, so its trace is $o_p(1)$ if and only if every entry is $o_p(1)$. Thus, we have

$$\frac{1}{N} \left[\Lambda'^* M' \hat{\Sigma}_{ee}^{-1} M \Lambda^* - \Lambda'^* M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} (\hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \Lambda^* \right] \xrightarrow{p} 0. \quad (\text{A.13})$$

Let $A \equiv (\hat{\Lambda} - \Lambda^*)' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}_N^{-1}$. Then $I_r - A = \Lambda'^* M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}_N^{-1}$. So equation (A.13) simplifies to

$$\frac{1}{N} \Lambda'^* M' \hat{\Sigma}_{ee}^{-1} M \Lambda^* - (I_r - A) \frac{1}{N} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} (I_r - A)' \xrightarrow{p} 0.$$

By Lemma A.2(a) and (A.9), we know $\frac{1}{N}\Lambda^*M'\hat{\Sigma}_{ee}^{-1}M\Lambda^* = \frac{1}{N}\Lambda^*M'\Sigma_{ee}^{*-1}M\Lambda^* + o_p(1)$. Thus,

$$\frac{1}{N}\Lambda^*M'\Sigma_{ee}^{*-1}M\Lambda^* - (I_r - A)\frac{1}{N}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}(I_r - A)' \xrightarrow{p} 0. \quad (\text{A.14})$$

By Assumption C.3, the expression $\frac{1}{N}\Lambda^*M'\Sigma_{ee}^{*-1}M\Lambda^*$ is positive definite in the limit, so the second term is of full rank in the limit which implies that $(I_r - A)$ is of full rank in the limit.

Alternatively, equation (A.13) can be rewritten as

$$\frac{1}{N}(\hat{\Lambda} - \Lambda^*)'M'\hat{\Sigma}_{ee}^{-1}M(\hat{\Lambda} - \Lambda^*) - A\left(\frac{1}{N}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}\right)A' \xrightarrow{p} 0. \quad (\text{A.15})$$

We now make use of the first-order conditions to proceed the proof. The first-order condition (3.3) post-multiplied by $\hat{\Lambda}$ implies

$$\hat{\Lambda}'M'\hat{\Sigma}_{zz}^{-1}(M_{zz} - \hat{\Sigma}_{zz})\hat{\Sigma}_{zz}^{-1}M\hat{\Lambda} = 0.$$

By (A.2), the above equation can be simplified as

$$\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}(M_{zz} - \hat{\Sigma}_{zz})\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda} = 0,$$

which is equivalent to

$$\begin{aligned} & \hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda} = -\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}(\hat{\Sigma}_{ee} - \Sigma_{ee}^*)\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda} \\ & + \hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}M\Lambda^*\Lambda^*M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda} + \hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}M\Lambda^*\frac{1}{T}\sum_{t=1}^T f_t^*e_t'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda} \\ & + \hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}\frac{1}{T}\sum_{t=1}^T e_t f_t^* \Lambda^*M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda} + \hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}\frac{1}{T}\sum_{t=1}^T (e_t e_t' - \Sigma_{ee}^*)\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}. \end{aligned}$$

With notations of \hat{P} and A , we have

$$\begin{aligned} I_r &= (I_r - A)'(I_r - A) + \frac{1}{N^2}\hat{P}^{-1}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}\frac{1}{T}\sum_{t=1}^T (e_t e_t' - \Sigma_{ee}^*)\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}\hat{P}^{-1} \\ & + (I_r - A)'\frac{1}{NT}\sum_{t=1}^T f_t^*e_t'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}\hat{P}^{-1} + \frac{1}{N}\hat{P}^{-1}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}\frac{1}{T}\sum_{t=1}^T e_t f_t^* (I_r - A) \\ & - \frac{1}{N^2}\hat{P}^{-1}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}(\hat{\Sigma}_{ee} - \Sigma_{ee}^*)\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}\hat{P}^{-1} = i_1 + i_2 + \dots + i_5, \quad \text{say} \end{aligned} \quad (\text{A.16})$$

Term i_2 is $\|\hat{P}^{-1/2}\|^2 \cdot O_p(T^{-1/2})$ by Lemma A.3(a). Term i_3 is $\|I - A\| \cdot \|\hat{P}^{-1/2}\| \cdot O_p(T^{-1/2})$ by Lemma A.3(b). Term i_4 is the transpose of i_3 and therefore has the same convergence rate as i_3 . The last term is $o_p(1)$ by Lemma A.3(c) and (A.12). Given these results, we have

$$I_r = (I - A)'(I - A) + \|\hat{P}^{-1/2}\|^2 O_p(T^{-1/2}) + \|I - A\| \cdot \|\hat{P}^{-1/2}\| \cdot O_p(T^{-1/2}) + o_p(1). \quad (\text{A.17})$$

Moreover, by the definition of \hat{P} , equation (A.14) yields

$$\left(\frac{1}{N}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}\right)^{-1} = (I_r - A)' \left(\frac{1}{N}\Lambda^*M'\Sigma_{ee}^{*-1}M\Lambda^*\right)^{-1} (I_r - A) + o_p(\|I_r - A\|^2).$$

This implies that

$$\|\hat{P}^{-1/2}\|^2 = \text{tr}(\hat{P}^{-1}) = \text{tr}\left[(I_r - A)' \left(\frac{1}{N}\Lambda^*M'\Sigma_{ee}^{*-1}M\Lambda^*\right)^{-1} (I_r - A) + o_p(\|I_r - A\|^2)\right].$$

The right hand side is at most $O_p[(A^2) \vee 1]$, implying that $\|\hat{P}^{-1/2}\| = O_p(A \vee 1)$, where $a \vee b$ denotes the maximum of a and b . So together with (A.17), we obtain $A = O_p(1)$. To see this, notice that the left hand side of equation (A.17) is bounded. Hence, if $A \neq O_p(1)$, then A is stochastically unbounded, the right hand side of (A.17) is dominated by $A'A$ in view of $\|\hat{P}^{-1/2}\| = O_p(A)$, but $A'A$ diverges. Then a contradiction arises. Thus, $A = O_p(1)$, which in turn implies that $\|\hat{P}^{-1/2}\| = O_p(1)$, or equivalently $\|\hat{P}^{-1}\| = O_p(1)$.

Now we sharpen the result to $A = o_p(1)$. From equation (A.17), $\|\hat{P}^{-1/2}\| = O_p(1)$ and $A = O_p(1)$, we have

$$(I_r - A)'(I_r - A) - I_r \xrightarrow{p} 0.$$

And from (A.14),

$$\frac{1}{N}\Lambda^*M'\Sigma_{ee}^{*-1}M\Lambda^* - (I_r - A)\frac{1}{N}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}(I_r - A)' = o_p(1).$$

By the identification condition, $\frac{1}{N}\Lambda^*M'\Sigma_{ee}^{*-1}M\Lambda^*$ and $\frac{1}{N}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}$ are both diagonal with distinct diagonal elements. Applying Lemma A.1 of the supplement of Bai and Li (2012) to the preceding two equations, we have that $I_r - A$ converges in probability to a diagonal matrix with diagonal elements either 1 or -1. By correctly choosing the column signs, the case -1 is precluded. Therefore, we have $I_r - A \xrightarrow{p} I_r$, or equivalently $A = o_p(1)$.

Next, we consider the first-order condition on Λ (equation (3.3)). By (A.2), we can simplify equation (3.3) as

$$\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}(M_{zz} - \hat{\Sigma}_{zz})\hat{\Sigma}_{ee}^{-1}M = 0.$$

Using the expression of M_{zz} , we can write the preceding equation as

$$\begin{aligned} \hat{\Lambda}' - \Lambda^{*'} &= -A'\Lambda^{*'} + (I - A)'\frac{1}{T}\sum_{t=1}^T f_t^* e_t' \hat{\Sigma}_{ee}^{-1} M \hat{R}_N^{-1} + \hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T e_t f_t^{*'} \Lambda^{*'} \quad (\text{A.18}) \\ &+ \hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T [e_t e_t' - \Sigma_{ee}^*] \hat{\Sigma}_{ee}^{-1} M \hat{R}_N^{-1} - \hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} (\hat{\Sigma}_{ee} - \Sigma_{ee}^*) \hat{\Sigma}_{ee}^{-1} M \hat{R}_N^{-1}. \end{aligned}$$

By $A = o_p(1)$ and Lemma A.3 (d), we have that the first two terms are $o_p(1)$. By $\|\hat{P}^{-1}\| = O_p(1)$ and Lemma A.3 (b), the third term is $o_p(1)$. By $\|\hat{P}^{-1}\| = O_p(1)$ and Lemma A.3 (e), the fourth term is $o_p(1)$. By $\|\hat{P}^{-1}\| = O_p(1)$ and Lemma A.3 (f), the last term is $o_p(1)$. Given the above result, we have $\hat{\Lambda}' - \Lambda^{*'} = o_p(1)$, which implies that $\hat{\Lambda} \xrightarrow{p} \Lambda^{*}$. This completes the proof of Proposition 4.1. \square

Corollary A.1 *Under Assumptions A-D,*

- (a) $\frac{1}{N}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda} - \frac{1}{N}\Lambda^*M'\Sigma_{ee}^{*-1}M\Lambda^* = o_p(1);$
- (b) $\hat{P}_N = O_p(N), \hat{P} = O_p(1), \hat{G} = O_p(N^{-1}), \hat{G}_N = O_p(1);$
- (c) $\frac{1}{N}(\hat{\Lambda} - \Lambda)'M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda} = o_p(1).$

PROOF OF COROLLARY A.1. Result (a) follows from equation (A.14) and $A = (\hat{\Lambda} - \Lambda)'M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}\hat{P}_N^{-1} = o_p(1).$

For part (b), by Assumption C.3, $N^{-1}\Lambda^*M'\Sigma_{ee}^{*-1}M\Lambda^* \rightarrow P_\infty > 0.$ This result, together with result (a) of this corollary, implies $\hat{P} = O_p(1)$ and therefore $\hat{P}_N = O_p(N).$ From $\hat{G} = (I_r + \hat{P}_N)^{-1},$ we have $\hat{G} = O_p(N^{-1})$ and hence $\hat{G}_N = O_p(1).$

Result (c) follows from $\hat{P} = O_p(N)$ and $A = o_p(1).$ \square

Appendix B: Proofs of Theorems 4.1, 4.2 and 4.5

Hereafter, for notational simplicity, we drop “*” from the symbols of underlying true values. The following lemmas are used in the proofs of Theorems 4.1 and 4.2.

Lemma B.1 *Under Assumptions A-D,*

- (a) $\hat{P}_N^{-1}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}\frac{1}{T}\sum_{t=1}^T(e_t e_t' - \Sigma_{ee})\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}\hat{P}_N^{-1} = O_p(T^{-1/2});$
- (b) $\hat{P}_N^{-1}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}\frac{1}{T}\sum_{t=1}^T e_t f_t' = O_p(T^{-1/2});$
- (c) $\hat{P}_N^{-1}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}(\hat{\Sigma}_{ee} - \Sigma_{ee})\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}\hat{P}_N^{-1} = \frac{1}{\sqrt{N}}O_p\left(\left[\frac{1}{N}\sum_{i=1}^N(\hat{\sigma}_i^2 - \sigma_i^2)^2\right]^{\frac{1}{2}}\right);$
- (d) $\frac{1}{T}\sum_{t=1}^T f_t e_t' \hat{\Sigma}_{ee}^{-1}M\hat{R}_N^{-1} = O_p(T^{-1/2});$
- (e) $\hat{P}_N^{-1}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}\frac{1}{T}\sum_{t=1}^T [e_t e_t' - \Sigma_{ee}]\hat{\Sigma}_{ee}^{-1}M\hat{R}_N^{-1} = O_p(T^{-1/2});$
- (f) $\hat{P}_N^{-1}\hat{\Lambda}'M'\hat{\Sigma}_{ee}^{-1}(\hat{\Sigma}_{ee} - \Sigma_{ee})\hat{\Sigma}_{ee}^{-1}M\hat{R}_N^{-1} = \frac{1}{\sqrt{N}}O_p\left(\left[\frac{1}{N}\sum_{i=1}^N(\hat{\sigma}_i^2 - \sigma_i^2)^2\right]^{\frac{1}{2}}\right).$

PROOF OF LEMMA B.1. First, we consider (a). The left hand side is equal to

$$\hat{P}^{-1}\frac{1}{N^2}\left[\sum_{i=1}^N\sum_{j=1}^N\left(\sum_{p=1}^k\hat{\lambda}_p m_{ip}\right)\frac{1}{\hat{\sigma}_i^2\hat{\sigma}_j^2}\frac{1}{T}\sum_{t=1}^T[e_{it}e_{jt} - E(e_{it}e_{jt})]\left(\sum_{q=1}^k m_{jq}\hat{\lambda}'_q\right)\right]\hat{P}^{-1},$$

which is bounded in norm by

$$C^2\|\hat{P}^{-1}\|^2\left[\frac{1}{N}\sum_{i=1}^N\frac{1}{\hat{\sigma}_i^2}\left\|\sum_{p=1}^k\hat{\lambda}_p m_{ip}\right\|^2\right]\left[\frac{1}{N^2}\sum_{i=1}^N\sum_{j=1}^N\left|\frac{1}{T}\sum_{t=1}^T[e_{it}e_{jt} - E(e_{it}e_{jt})]\right|^2\right]^{1/2}.$$

Moreover, by Corollary A.1(a), we have

$$\frac{1}{N} \sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2} \left\| \sum_{p=1}^k \hat{\lambda}_p m_{ip} \right\|^2 = \text{tr} \left[\frac{1}{N} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \right] \xrightarrow{p} \text{tr} \left[\frac{1}{N} \Lambda' M' \Sigma_{ee}^{-1} M \Lambda \right] = \text{tr}(P). \quad (\text{B.1})$$

By

$$E \left[\frac{1}{N^2} \sum_{i=1}^N \sum_{j=1}^N \left| \frac{1}{T} \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})] \right|^2 \right] = O(T^{-1}),$$

together with Corollary A.1(b) and (B.1), we obtain (a).

Next, we consider (b). The left hand side can be written as

$$\hat{P}^{-1} \frac{1}{N} \sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2} \left(\sum_{p=1}^k \hat{\lambda}_p m_{ip} \right) \frac{1}{T} \sum_{t=1}^T e_{it} f'_t,$$

which is bounded in norm by

$$C \|\hat{P}^{-1}\| \left[\frac{1}{N} \sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2} \left\| \sum_{p=1}^k \hat{\lambda}_p m_{ip} \right\|^2 \right]^{1/2} \left[\frac{1}{N} \sum_{i=1}^N \left\| \frac{1}{T} \sum_{t=1}^T e_{it} f'_t \right\|^2 \right]^{1/2},$$

which is $O_p(T^{-1/2})$ by (B.1). Thus, (b) follows.

For part (c), the left hand side can be written as

$$\hat{P}_N^{-1/2} \left[\sum_{i=1}^N \hat{P}_N^{-1/2} \left(\sum_{p=1}^k \hat{\lambda}_p m_{ip} \right) \frac{\hat{\sigma}_i^2 - \sigma_i^2}{\hat{\sigma}_i^4} \left(\sum_{q=1}^k m_{iq} \hat{\lambda}'_q \right) \hat{P}_N^{-1/2} \right] \hat{P}_N^{-1/2},$$

which is bounded in norm by

$$C^2 \|\hat{P}_N^{-1/2}\|^2 \cdot \sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2} \left\| \hat{P}_N^{-1/2} \left(\sum_{p=1}^k \hat{\lambda}_p m_{ip} \right) \right\|^2 (\hat{\sigma}_i^2 - \sigma_i^2). \quad (\text{B.2})$$

Since

$$\sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2} \left\| \hat{P}_N^{-1/2} \sum_{p=1}^k \hat{\lambda}_p m_{ip} \right\|^2 = r$$

by (A.5), this gives

$$\frac{1}{\hat{\sigma}_i} \left\| \hat{P}_N^{-1/2} \sum_{p=1}^k \hat{\lambda}_p m_{ip} \right\| \leq \sqrt{r}.$$

Hence, expression in (B.2) is bounded by

$$C^2 \sqrt{r} \|\hat{P}_N^{-1/2}\|^2 \cdot \sum_{i=1}^N \frac{1}{\hat{\sigma}_i} \left\| \hat{P}_N^{-1/2} \left(\sum_{p=1}^k \hat{\lambda}_p m_{ip} \right) \right\| (\hat{\sigma}_i^2 - \sigma_i^2),$$

which is further bounded by

$$C^2 \sqrt{r} \|\hat{P}_N^{-1/2}\|^2 \cdot \left[\sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2} \left\| \hat{P}_N^{-1/2} \sum_{p=1}^k \hat{\lambda}_p m_{ip} \right\|^2 \right]^{1/2} \left[\sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2 \right]^{1/2}.$$

Then result (c) follows by noticing that $\hat{P}_N = O_p(N)$.

The proofs of the remaining three parts are similar to those of the first three. The details are therefore omitted. \square

Lemma B.2 *Under Assumptions A-D,*

$$A \equiv (\hat{\Lambda} - \Lambda)' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}_N^{-1} = O_p(T^{-1/2}) + O_p(\|\hat{\Lambda} - \Lambda\|^2) + O_p\left(\left[\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2\right]^{\frac{1}{2}}\right).$$

PROOF OF LEMMA B.2. Consider equation (A.16) in the proof of Proposition 4.1, we had shown $A = o_p(1)$. So term AA' is of a smaller order and hence negligible. With Lemma B.2 (a), (b) and (c), equation (A.16) can be simplified as

$$A + A' = O_p(T^{-1/2}) + o_p\left(\left[\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2\right]^{\frac{1}{2}}\right). \quad (\text{B.3})$$

By the identification condition, we know both $\Lambda' (\frac{1}{N} M' \Sigma_{ee}^{-1} M) \Lambda$ and $\hat{\Lambda}' (\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M) \hat{\Lambda}$ are diagonal matrices, which implies

$$\text{Ndg} \left\{ \Lambda' \left(\frac{1}{N} M' \Sigma_{ee}^{-1} M \right) \Lambda - \hat{\Lambda}' \left(\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M \right) \hat{\Lambda} \right\} = 0,$$

where Ndg denotes the operator which sets the diagonal elements of its input to zeros. By adding and subtracting terms,

$$\begin{aligned} & \text{Ndg} \left\{ (\hat{\Lambda} - \Lambda)' \left(\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M \right) \hat{\Lambda} + \hat{\Lambda}' \left(\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M \right) (\hat{\Lambda} - \Lambda) \right. \\ & \left. - (\hat{\Lambda} - \Lambda)' \left(\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M \right) (\hat{\Lambda} - \Lambda) + \Lambda' \left[\frac{1}{N} M' (\hat{\Sigma}_{ee}^{-1} - \Sigma_{ee}^{-1}) M \right] \Lambda \right\} = 0. \end{aligned} \quad (\text{B.4})$$

By Lemma A.2 (b), $\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M = \frac{1}{N} M' \Sigma_{ee}^{-1} M + o_p(1) = R + o_p(1)$, where the last equation is due to Assumption C.3. So term $(\hat{\Lambda} - \Lambda)' (\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M) (\hat{\Lambda} - \Lambda) = O_p(\|\hat{\Lambda} - \Lambda\|^2)$. Given this result, together with Lemma A.2(a), we have

$$\begin{aligned} & \text{Ndg} \left\{ (\hat{\Lambda} - \Lambda)' \left(\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M \right) \hat{\Lambda} + \hat{\Lambda}' \left(\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M \right) (\hat{\Lambda} - \Lambda) \right\} \\ & = O_p(\|\hat{\Lambda} - \Lambda\|^2) + O_p\left(\left[\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2\right]^{1/2}\right). \end{aligned} \quad (\text{B.5})$$

Notice that $(\hat{\Lambda} - \Lambda)' (\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M) \hat{\Lambda} = (\hat{\Lambda} - \Lambda)' (\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M) \hat{\Lambda} \hat{P}^{-1} \hat{P} = A \hat{P}$, where the last inequality is due to the definition of A . By $\hat{P} = P + o_p(1)$ from Corollary A.1 (a), we have

$$(\hat{\Lambda} - \Lambda)' \left(\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M \right) \hat{\Lambda} = AP + o_p(A).$$

According to the preceding result, we can rewrite (B.5) as

$$\text{Ndg}\{AP + PA'\} = O_p(\|\hat{\Lambda} - \Lambda\|^2) + O_p\left(\left[\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2\right]^{1/2}\right), \quad (\text{B.6})$$

where $o_p(A)$ is discarded since it has an smaller order than other terms.

Now equation (B.3) has $\frac{1}{2}r(r+1)$ restrictions and equation (B.6) has $\frac{1}{2}r(r-1)$ restrictions, the $r \times r$ matrix A can be uniquely determined. Solving this linear equation system, we have

$$A = O_p(T^{-1/2}) + O_p(\|\hat{\Lambda} - \Lambda\|^2) + O_p\left(\left[\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2\right]^{\frac{1}{2}}\right).$$

This completes the proof. \square

PROOF OF THEOREM 4.1. We first consider the first order condition (3.4), which can be written as

$$\text{diag} \left\{ (M_{zz} - \hat{\Sigma}_{zz}) - (M_{zz} - \hat{\Sigma}_{zz}) \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{G} \hat{\Lambda}' M' - M \hat{\Lambda} \hat{G} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} (M_{zz} - \hat{\Sigma}_{zz}) \right\} = 0,$$

where ‘‘diag’’ denotes the diagonal operator and $\hat{G} = (I_r + \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1}$. By

$$M_{zz} = M \Lambda \Lambda' M' + \Sigma_{ee} + M \Lambda \frac{1}{T} \sum_{t=1}^T f_t e_t' + \frac{1}{T} \sum_{t=1}^T e_t f_t' \Lambda' M' + \frac{1}{T} \sum_{t=1}^T (e_t e_t' - \Sigma_{ee}),$$

with some algebra manipulations, we can further write the preceding equation as

$$\begin{aligned} \hat{\sigma}_i^2 - \sigma_i^2 &= \frac{1}{T} \sum_{t=1}^T (e_{it}^2 - \sigma_i^2 + 2m_i' \Lambda \frac{1}{T} \sum_{t=1}^T f_t e_{it} - 2m_i' \hat{\Lambda} \hat{G} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \Lambda \frac{1}{T} \sum_{t=1}^T f_t e_{it} \\ &\quad - 2m_i' \Lambda \frac{1}{T} \sum_{t=1}^T f_t e_t' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{G} \hat{\Lambda}' m_i - 2m_i' \hat{\Lambda} \hat{G} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T [e_t e_{it} - E(e_t e_{it})]) \quad (\text{B.7}) \\ &+ m_i' (\hat{\Lambda} - \Lambda) (\hat{\Lambda} - \Lambda)' m_i - 2m_i' (\hat{\Lambda} - \Lambda) \hat{\Lambda}' m_i + 2m_i' (\hat{\Lambda} - \Lambda) \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{G} \hat{\Lambda}' m_i \\ &\quad + 2m_i' \Lambda (\hat{\Lambda} - \Lambda)' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{G} \hat{\Lambda}' m_i + 2 \frac{\hat{\sigma}_i^2 - \sigma_i^2}{\hat{\sigma}_i^2} m_i' \hat{\Lambda} \hat{G} \hat{\Lambda}' m_i. \end{aligned}$$

By $\hat{G} \hat{P}_N = \hat{P}_N \hat{G} = I_N - \hat{G}$, we have $\hat{G} = (I_N - \hat{G}) \hat{P}_N^{-1} = \hat{P}_N^{-1} (I_N - \hat{G})$. Then, the third term on right hand side (ignoring the factor 2) is equal to

$$m_i' \hat{\Lambda} (I_N - \hat{G}) \hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \Lambda \frac{1}{T} \sum_{t=1}^T f_t e_{it} = m_i' \hat{\Lambda} (I_N - \hat{G}) (I - A)' \frac{1}{T} \sum_{t=1}^T f_t e_{it} \quad (\text{B.8})$$

and the sum of the seventh and eighth terms is equal to $-2m_i' (\hat{\Lambda} - \Lambda) \hat{G} \hat{\Lambda}' m_i$. Define

$$\check{\psi} = \frac{1}{T} \sum_{t=1}^T f_t e_t' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}_N^{-1}; \quad \check{\phi} = \hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T (e_t e_t' - \Sigma_{ee}) \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}_N^{-1}.$$

Now consider the sum of the fourth and ninth terms. By $\hat{G} = \hat{P}_N^{-1} (I_N - \hat{G})$, together with the definitions of $\check{\psi}$, this term is equal to

$$\begin{aligned} &-2m_i' \Lambda \frac{1}{T} \sum_{t=1}^T f_t e_t' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{G} \hat{\Lambda}' m_i + 2m_i' \Lambda (\hat{\Lambda} - \Lambda)' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{G} \hat{\Lambda}' m_i \\ &= -2m_i' \Lambda \check{\psi} (I_N - \hat{G}) \hat{\Lambda}' m_i + 2m_i' \Lambda A (I_N - \hat{G}) \hat{\Lambda}' m_i \end{aligned}$$

$$\begin{aligned}
&= 2m'_i \Lambda \ddot{\psi} \hat{G} \hat{\Lambda}' m_i - 2m'_i \Lambda A \hat{G} \hat{\Lambda}' m_i - 2m'_i \Lambda \ddot{\psi} (\hat{\Lambda} - \Lambda)' m_i + 2m'_i \Lambda A (\hat{\Lambda} - \Lambda)' m_i \\
&\quad + m'_i \Lambda (A + A' - \ddot{\psi} - \ddot{\psi}') \Lambda' m_i.
\end{aligned}$$

Also, by (A.16), we have

$$A' + A = A' A + \ddot{\phi} + (I_r - A)' \ddot{\psi} + \ddot{\psi}' (I_r - A) - \hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} (\hat{\Sigma}_{ee} - \Sigma_{ee}) \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}_N^{-1},$$

or equivalently

$$A' + A - \ddot{\psi} - \ddot{\psi}' = A' A + \ddot{\phi} - A' \ddot{\psi} - \ddot{\psi}' A - \hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} (\hat{\Sigma}_{ee} - \Sigma_{ee}) \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}_N^{-1}.$$

Thus, it follows that

$$\begin{aligned}
&-2m'_i \Lambda \frac{1}{T} \sum_{t=1}^T f_t e_t' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{G} \hat{\Lambda}' m_i + 2m'_i \Lambda (\hat{\Lambda} - \Lambda)' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{G} \hat{\Lambda}' m_i \quad (\text{B.9}) \\
&= 2m'_i \Lambda \ddot{\psi} \hat{G} \hat{\Lambda}' m_i - 2m'_i \Lambda A \hat{G} \hat{\Lambda}' m_i - 2m'_i \Lambda \ddot{\psi} (\hat{\Lambda} - \Lambda)' m_i + 2m'_i \Lambda A (\hat{\Lambda} - \Lambda)' m_i - m'_i \Lambda A' A \Lambda' m_i \\
&\quad - m'_i \Lambda \ddot{\phi} \Lambda' m_i + 2m'_i \Lambda A' \ddot{\psi} \Lambda' m_i + m'_i \Lambda \hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} (\hat{\Sigma}_{ee} - \Sigma_{ee}) \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}_N^{-1} \Lambda' m_i.
\end{aligned}$$

Using (B.8) and (B.9), we can rewrite (B.7) as

$$\begin{aligned}
\hat{\sigma}_i^2 - \sigma_i^2 &= \frac{1}{T} \sum_{t=1}^T (e_{it}^2 - \sigma_i^2) - 2m'_i (\hat{\Lambda} - \Lambda) \frac{1}{T} \sum_{t=1}^T f_t e_{it} + 2m'_i \hat{\Lambda} \hat{G} \frac{1}{T} \sum_{t=1}^T f_t e_{it} \quad (\text{B.10}) \\
&\quad + 2m'_i \hat{\Lambda} A' \frac{1}{T} \sum_{t=1}^T f_t e_{it} - 2m'_i \hat{\Lambda} \hat{G} A' \frac{1}{T} \sum_{t=1}^T f_t e_{it} + 2m'_i \Lambda \ddot{\psi} \hat{G} \hat{\Lambda}' m_i \\
&\quad - 2m'_i \Lambda A \hat{G} \hat{\Lambda}' m_i - 2m'_i \Lambda \ddot{\psi} (\hat{\Lambda} - \Lambda)' m_i + 2m'_i \Lambda A (\hat{\Lambda} - \Lambda)' m_i \\
&\quad + m'_i \Lambda A' A \Lambda' m_i - 2m'_i \Lambda A' \ddot{\psi} \Lambda' m_i - 2m'_i (\hat{\Lambda} - \Lambda) \hat{G} \hat{\Lambda}' m_i + 2 \frac{\hat{\sigma}_i^2 - \sigma_i^2}{\hat{\sigma}_i^2} m'_i \hat{\Lambda} \hat{G} \hat{\Lambda}' m_i \\
&\quad + m'_i \Lambda \ddot{\phi} \Lambda' m_i - m'_i \Lambda \hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} (\hat{\Sigma}_{ee} - \Sigma_{ee}) \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}_N^{-1} \Lambda' m_i \\
&\quad - 2m'_i \hat{\Lambda} \hat{G} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T [e_t e_{it} - E(e_t e_{it})] + m'_i (\hat{\Lambda} - \Lambda) (\hat{\Lambda} - \Lambda)' m_i \\
&= a_{i,1} + a_{i,2} + \cdots + a_{i,17}, \quad \text{say.}
\end{aligned}$$

By the Cauchy-Schwartz inequality, we have

$$\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2 \leq 17 \frac{1}{N} \sum_{i=1}^N (\|a_{i,1}\|^2 + \cdots + \|a_{i,17}\|^2).$$

The first term $N^{-1} \sum_{i=1}^N \|a_{i,1}\|^2 = O_p(T^{-1})$ by

$$E \left[\frac{1}{N} \sum_{i=1}^N \left| \frac{1}{T} \sum_{t=1}^T (e_{it}^2 - \sigma_i^2) \right|^2 \right] = O(T^{-1}).$$

The second term is bounded in norm by

$$4C^2 \|\hat{\Lambda} - \Lambda\|^2 \frac{1}{N} \sum_{i=1}^N \left\| \frac{1}{T} \sum_{t=1}^T f_t e_{it} \right\|^2 = o_p(T^{-1})$$

by $\hat{\Lambda} - \Lambda = o_p(1)$ and

$$E \left[\frac{1}{N} \sum_{i=1}^N \left\| \frac{1}{T} \sum_{t=1}^T f_t e_{it} \right\|^2 \right] = O(T^{-1}).$$

Similarly, one can show that the 3rd, 4th, 5th, 6th, 8th, 11th and 14th terms are all $o_p(T^{-1})$. The 7th term is bounded in norm by

$$(4\|\Lambda\|^2 \cdot \|\hat{\Lambda}\|^2 \cdot \|\hat{G}\|^2 \cdot \|A\|^2) \frac{1}{N} \sum_{i=1}^N \|m_i\|^4,$$

which is $O_p(N^{-2}T^{-1}) + O_p(N^{-2}) \cdot O_p(\|\hat{\Lambda} - \Lambda\|^4) + O_p(N^{-2}) \cdot O_p[\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)]$ by $\hat{G} = O_p(N^{-1})$, $\hat{\Lambda} = \Lambda + o_p(1)$ and Lemma B.2. This result can be simplified to $\frac{1}{N} \sum_{i=1}^N \|a_{i,7}\|^2 = o_p(T^{-1}) + o_p(\|\hat{\Lambda} - \Lambda\|^2)$ since $O_p(N^{-2}) \cdot O_p[\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)]$ is of smaller order than $\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2$. Similar to the 7th term, the 9th and 10th terms are both of the order $o_p(T^{-1}) + o_p(\|\hat{\Lambda} - \Lambda\|^2)$. The 12th term is $o_p(\|\hat{\Lambda} - \Lambda\|^2)$ by $\hat{G} = O_p(N^{-1})$. The 13th term is of smaller order term than $\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)$ and therefore negligible. The 15th term is $o_p(\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2))$ by Lemma B.1 (f). The 16th term is $O_p(T^{-1})$. The last term is $O_p(\|\hat{\Lambda} - \Lambda\|^4)$. Given the above results, we have

$$\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2 = O_p(T^{-1}) + o_p(\|\hat{\Lambda} - \Lambda\|^2). \quad (\text{B.11})$$

Next, we derive bounds for $\|\hat{\Lambda} - \Lambda\|^2$. By equation (A.18), together with Lemma B.1(b), (d), (e) and (f) and Lemma B.2, we have

$$\hat{\Lambda} - \Lambda = O_p(T^{-1/2}) + O_p\left(\left[\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2\right]^{1/2}\right). \quad (\text{B.12})$$

Substituting equation (B.12) into (B.11), we have $\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2 = O_p(T^{-1})$. This proves the second result of Theorem 4.1. \square

To prove the first result of Theorem 4.1, we need the following lemmas.

Lemma B.3 *Under Assumptions A-D, we have*

- (a) $\hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T (e_t e_t' - \Sigma_{ee}) \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}_N^{-1}$
 $= O_p(N^{-1}T^{-1/2}) + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2});$
- (b) $\hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T e_t f_t' = O_p(N^{-1/2}T^{-1/2}) + O_p(T^{-1});$
- (c) $\hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} (\hat{\Sigma}_{ee} - \Sigma_{ee}) \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}_N^{-1} = O_p(N^{-1}T^{-1/2});$
- (d) $\frac{1}{T} \sum_{t=1}^T f_t e_t' \hat{\Sigma}_{ee}^{-1} M \hat{R}_N^{-1} = O_p(N^{-1/2}T^{-1/2}) + O_p(T^{-1});$
- (e) $\hat{P}_N^{-1} \hat{\Lambda}' \left(M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T [e_t e_t' - \Sigma_{ee}] \hat{\Sigma}_{ee}^{-1} M \right) \hat{R}_N^{-1}$

$$= O_p(N^{-1}T^{-1/2}) + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2});$$

$$(f) \hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} (\hat{\Sigma}_{ee} - \Sigma_{ee}) \hat{\Sigma}_{ee}^{-1} M \hat{R}_N^{-1} = O_p(N^{-1}T^{-1/2}).$$

PROOF OF LEMMA B.3. We first consider (a). We rewrite it as

$$\hat{P}^{-1} \hat{\Lambda}' \left(\frac{1}{N^2} M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T (e_t e_t' - \Sigma_{ee}) \hat{\Sigma}_{ee}^{-1} M \right) \hat{\Lambda} \hat{P}^{-1}.$$

Since we already know that $\|\hat{P}^{-1}\| = O_p(1)$ and $\|\hat{\Lambda}'\| = O_p(1)$, we only need to consider the term in the big parenthesis, which is

$$\begin{aligned} & \frac{1}{N^2 T} \sum_{i=1}^N \sum_{j=1}^N m_i m_j' \frac{1}{\hat{\sigma}_i^2 \hat{\sigma}_j^2} \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})] \\ &= \frac{1}{N^2 T} \sum_{i=1}^N \sum_{j=1}^N m_i m_j' \left(\frac{1}{\hat{\sigma}_i^2} - \frac{1}{\sigma_i^2} \right) \left(\frac{1}{\hat{\sigma}_j^2} - \frac{1}{\sigma_j^2} \right) \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})] \\ & \quad + \frac{1}{N^2 T} \sum_{i=1}^N \sum_{j=1}^N m_i m_j' \frac{1}{\sigma_i^2} \left(\frac{1}{\hat{\sigma}_j^2} - \frac{1}{\sigma_j^2} \right) \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})] \\ & \quad + \frac{1}{N^2 T} \sum_{i=1}^N \sum_{j=1}^N m_i m_j' \frac{1}{\sigma_j^2} \left(\frac{1}{\hat{\sigma}_i^2} - \frac{1}{\sigma_i^2} \right) \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})] \\ & \quad + \frac{1}{N^2 T} \sum_{i=1}^N \sum_{j=1}^N m_i m_j' \frac{1}{\sigma_i^2 \sigma_j^2} \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})]. \end{aligned}$$

By the Cauchy-Schwarz inequality, one can show the first term is bounded in norm by

$$C^8 \left(\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2 \right) \left(\frac{1}{N^2} \sum_{i=1}^N \sum_{j=1}^N \left\| \frac{1}{T} \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})] \right\|^2 \right)^{1/2},$$

which is $O_p(T^{-3/2})$ by the second part of Theorem 4.1. The second term equals to

$$\begin{aligned} & \frac{1}{N^2 T} \sum_{i=1}^N \sum_{j=1}^N m_i m_j' \frac{1}{\sigma_i^2} \left(\frac{1}{\hat{\sigma}_j^2} - \frac{1}{\sigma_j^2} \right) \sum_{t=1}^T [e_{it} e_{jt} - E(e_{it} e_{jt})] \\ &= \frac{1}{N} \sum_{j=1}^N m_j' \left(\frac{1}{\hat{\sigma}_j^2} - \frac{1}{\sigma_j^2} \right) \left(\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} m_i [e_{it} e_{jt} - E(e_{it} e_{jt})] \right), \end{aligned}$$

which is bounded in norm by

$$C^4 \left[\frac{1}{N} \sum_{j=1}^N (\hat{\sigma}_j^2 - \sigma_j^2)^2 \right]^{1/2} \left[\frac{1}{N} \sum_{j=1}^N \left(\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} m_i [e_{it} e_{jt} - E(e_{it} e_{jt})] \right)^2 \right]^{1/2},$$

which is $O_p(N^{-1/2}T^{-1})$. Similarly, the third term is also $O_p(N^{-1/2}T^{-1})$. The last term is $O_p(N^{-1}T^{-1/2})$. Hence result (a) follows.

Next, we consider (b). The left hand side of (b) is equivalent to

$$\hat{P}^{-1} \hat{\Lambda}' \left(\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T e_t f_t' \right).$$

Similarly to (a), it suffices to consider the term inside the parenthesis, which is

$$\begin{aligned} \frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T e_t f_t' &= \frac{1}{N} \sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2} m_i \frac{1}{T} \sum_{t=1}^T e_{it} f_t' \\ &= \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} m_i f_t' e_{it} + \frac{1}{N} \sum_{i=1}^N \left(\frac{1}{\hat{\sigma}_i^2} - \frac{1}{\sigma_i^2} \right) \frac{1}{T} \sum_{t=1}^T m_i f_t' e_{it}. \end{aligned}$$

The first term is $O_p(N^{-1/2}T^{-1/2})$. The second term is bounded in norm by

$$C^4 \left[\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2 \right]^{1/2} \left[\frac{1}{N} \sum_{i=1}^N \left\| \frac{1}{T} \sum_{t=1}^T f_t e_{it} \right\|^2 \right]^{1/2},$$

which is $O_p(T^{-1})$ by the second part of Theorem 4.1. Hence, result (b) follows.

For part (c), the left hand side of (c) is equivalent to

$$\hat{P}^{-1} \hat{\Lambda}' \left(\frac{1}{N^2} M' \hat{\Sigma}_{ee}^{-1} (\hat{\Sigma}_{ee} - \Sigma_{ee}) \hat{\Sigma}_{ee}^{-1} M \right) \hat{\Lambda} \hat{P}^{-1}.$$

It suffices to consider the expression in the parenthesis:

$$\frac{1}{N^2} \sum_{i=1}^N m_i m_i' \frac{\hat{\sigma}_i^2 - \sigma_i^2}{\hat{\sigma}_i^4} \leq \frac{1}{N} \left(\frac{1}{N} \sum_{i=1}^N \|m_i\|^2 \right)^{1/2} \left(\frac{1}{N} \sum_{i=1}^N \|m_i'\|^2 \frac{(\hat{\sigma}_i^2 - \sigma_i^2)^2}{\hat{\sigma}_i^8} \right)^{1/2},$$

which is $O_p(N^{-1}T^{-1/2})$ by the second part of Theorem 4.1. This proves result (c). The proofs of results (d), (e) and (f) are similar to those of (a), (b) and (c). The details are therefore omitted. \square

Lemma B.4 *Under Assumptions A-D,*

$$A \equiv (\hat{\Lambda} - \Lambda)' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}_N^{-1} = O_p \left(\frac{1}{\sqrt{NT}} \right) + O_p \left(\frac{1}{T} \right) + O_p(\|\hat{\Lambda} - \Lambda\|^2).$$

PROOF OF LEMMA B.4. Consider equation (A.16). Using the results in Lemma B.3 and the fact that $A'A$ has an order smaller than that of A and is therefore negligible, we have

$$A + A' = O_p \left(\frac{1}{\sqrt{NT}} \right) + O_p \left(\frac{1}{T} \right). \quad (\text{B.13})$$

Now consider the term $\frac{1}{N} \Lambda' M' (\hat{\Sigma}_{ee}^{-1} - \Sigma_{ee}^{-1}) M \Lambda$, which can be written as

$$\begin{aligned} \frac{1}{N} \Lambda' M' (\hat{\Sigma}_{ee}^{-1} - \Sigma_{ee}^{-1}) M \Lambda &= -\Lambda' \left[\frac{1}{N} \sum_{i=1}^N m_i m_i' \frac{\hat{\sigma}_i^2 - \sigma_i^2}{\hat{\sigma}_i^2 \sigma_i^2} \right] \Lambda \\ &= -\Lambda' \left[\frac{1}{N} \sum_{i=1}^N m_i m_i' \frac{\hat{\sigma}_i^2 - \sigma_i^2}{\sigma_i^4} \right] \Lambda + \Lambda' \left[\frac{1}{N} \sum_{i=1}^N m_i m_i' \frac{(\hat{\sigma}_i^2 - \sigma_i^2)^2}{\hat{\sigma}_i^2 \sigma_i^4} \right] \Lambda. \end{aligned} \quad (\text{B.14})$$

The norm of the second expression on the right hand side of (B.14) is bounded by

$$C \frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2 = O_p(T^{-1}),$$

by the boundedness of $m_i, \hat{\sigma}_i^2, \sigma_i^2$ by Assumptions C and D. Substituting (B.10) into the first expression on the right hand side of (B.14) and using the same arguments as we did at before (B.11), one can show that the first expression is $O_p(\frac{1}{\sqrt{NT}}) + o_p(\frac{1}{T})$. Hence, we have

$$\frac{1}{N}\Lambda'M'(\hat{\Sigma}_{ee}^{-1} - \Sigma_{ee}^{-1})M\Lambda = O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T}\right). \quad (\text{B.15})$$

Now consider (B.4). Using the same arguments as in the derivation of (B.6) except that the result for $\frac{1}{N}\Lambda'M'(\hat{\Sigma}_{ee}^{-1} - \Sigma_{ee}^{-1})M\Lambda$ is given by (B.15) instead of $o_p([\frac{1}{N}\sum_{i=1}^N(\hat{\sigma}_i^2 - \sigma_i^2)^2]^{1/2})$, we have

$$\text{Ndg}\{AP + PA'\} = O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T}\right) + O_p(\|\hat{\Lambda} - \Lambda\|^2). \quad (\text{B.16})$$

Solving the equation system (B.13) and (B.16), we have

$$A = O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T}\right) + O_p(\|\hat{\Lambda} - \Lambda\|^2),$$

as asserted in this lemma. This proves Lemma B.4. \square

PROOF OF THEOREM 4.1 (CONTINUED). Using the results in Lemma B.3 and Lemma B.4 and noticing that $\|\hat{\Lambda} - \Lambda\|^2$ is of smaller order than $\hat{\Lambda} - \Lambda$ and therefore negligible, we have from (A.18)

$$\hat{\Lambda} - \Lambda = O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T}\right),$$

as asserted by the first result of Theorem 4.1. This completes the proof of Theorem 4.1.

Corollary B.1 *Under Assumptions A-D,*

$$A \equiv (\hat{\Lambda} - \Lambda)'M'\hat{\Sigma}_{ee}^{-1}M\hat{\Lambda}\hat{P}_N^{-1} = O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T}\right).$$

Corollary B.1 is a direct result of Lemma B.4 and Theorem 4.1.

Lemma B.5 *Under Assumptions A-D,*

- (a) $\frac{1}{T}\sum_{t=1}^T f_t e_t' \hat{\Sigma}_{ee}^{-1} M \hat{R}_N^{-1} = \frac{1}{T}\sum_{t=1}^T f_t e_t' \Sigma_{ee}^{-1} M R_N^{-1} + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2});$
- (b) $\hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T e_t f_t' = P_N^{-1} \Lambda' M' \Sigma_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T e_t f_t' + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2});$
- (c) $\frac{1}{N} M' (\hat{\Sigma}_{ee}^{-1} - \Sigma_{ee}^{-1}) M = -\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^4} m_i m_i' (e_{it}^2 - \sigma_i^2) + \frac{1}{NT} \sum_{i=1}^N m_i m_i' \frac{\kappa_{iA} - \sigma_i^4}{\sigma_i^4} + O_p(N^{-1}T^{-1/2}) + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2}).$

PROOF OF LEMMA B.5. Equation (B.10) can be written as

$$\hat{\sigma}_i^2 - \sigma_i^2 = \frac{1}{T} \sum_{t=1}^T (e_{it}^2 - \sigma_i^2) + \mathcal{R}_i, \quad (\text{B.17})$$

where

$$\mathcal{R}_i = -2m_i' \hat{\Lambda} \hat{G} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T [e_t e_{it} - E(e_t e_{it})] + \mathcal{S}_i$$

with

$$\begin{aligned} \mathcal{S}_i = & -2m_i' (\hat{\Lambda} - \Lambda) \frac{1}{T} \sum_{t=1}^T f_t e_{it} + 2m_i' \hat{\Lambda} \hat{G} \frac{1}{T} \sum_{t=1}^T f_t e_{it} + 2m_i' \hat{\Lambda} A' \frac{1}{T} \sum_{t=1}^T f_t e_{it} - 2m_i' \hat{\Lambda} \hat{G} A' \frac{1}{T} \sum_{t=1}^T f_t e_{it} \\ & + 2m_i' \Lambda \check{\psi} \hat{G} \hat{\Lambda}' m_i - 2m_i' \Lambda A \hat{G} \hat{\Lambda}' m_i - 2m_i' \Lambda \check{\psi} (\hat{\Lambda} - \Lambda)' m_i + 2m_i' \Lambda A (\hat{\Lambda} - \Lambda)' m_i \\ & + m_i' \Lambda A' A \Lambda' m_i - 2m_i' \Lambda A' \check{\psi} \Lambda' m_i - 2m_i' (\hat{\Lambda} - \Lambda) \hat{G} \hat{\Lambda}' m_i + 2 \frac{\hat{\sigma}_i^2 - \sigma_i^2}{\hat{\sigma}_i^2} m_i' \hat{\Lambda} \hat{G} \hat{\Lambda}' m_i \\ & + m_i' \Lambda \check{\phi} \Lambda' m_i - m_i' \Lambda \hat{P}_N^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} (\hat{\Sigma}_{ee} - \Sigma_{ee}) \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda} \hat{P}_N^{-1} \Lambda' m_i + m_i' (\hat{\Lambda} - \Lambda) (\hat{\Lambda} - \Lambda)' m_i. \end{aligned}$$

Given that $\check{\psi} = O_p(N^{-1/2}T^{-1/2}) + O_p(T^{-1})$ by Lemma B.3 (b), $\hat{\Lambda} - \Lambda = O_p(N^{-1/2}T^{-1/2}) + O_p(T^{-1})$ by Theorem 4.1, $A = O_p(N^{-1/2}T^{-1/2}) + O_p(T^{-1})$ by Corollary B.1, by the same arguments in the derivation of (B.10), we have

$$\frac{1}{N} \sum_{i=1}^N \mathcal{S}_i^2 = O_p(N^{-1}T^{-2}) + O_p(N^{-2}T^{-1}) + O_p(T^{-3}). \quad (\text{B.18})$$

We now consider

$$\frac{1}{N} \sum_{i=1}^N \left| m_i' \hat{\Lambda} \hat{G} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T [e_t e_{it} - E(e_t e_{it})] \right|^2,$$

which is bounded in norm by

$$C^2 \|\hat{\Lambda}\|^4 \cdot \|\hat{G}_N\|^2 \cdot \frac{1}{N} \sum_{i=1}^N \left| \frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T [e_t e_{it} - E(e_t e_{it})] \right|^2.$$

Since $\hat{\Lambda} = \Lambda + o_p(1)$ and $\hat{G}_N = O_p(1)$, it suffices to consider the term

$$\frac{1}{N} \sum_{i=1}^N \left| \frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T [e_t e_{it} - E(e_t e_{it})] \right|^2,$$

which, by the Cauchy-Schwarz inequality, is bounded by

$$\begin{aligned} & 2 \frac{1}{N} \sum_{i=1}^N \left| \frac{1}{NT} \sum_{j=1}^N \frac{1}{\sigma_j^2} m_j \sum_{t=1}^T [e_{jt} e_{it} - E(e_{jt} e_{it})] \right|^2 \\ & + 2 \frac{1}{N} \sum_{i=1}^N \left| \frac{1}{NT} \sum_{j=1}^N \frac{\hat{\sigma}_j^2 - \sigma_j^2}{\hat{\sigma}_j^2 \sigma_j^2} m_j \sum_{t=1}^T [e_{jt} e_{it} - E(e_{jt} e_{it})] \right|^2. \end{aligned}$$

The first expression is $O_p(N^{-1}T^{-1})$. The second expression is bounded by

$$C^{10} \left[\frac{1}{N} \sum_{j=1}^N (\hat{\sigma}_j^2 - \sigma_j^2)^2 \right] \left[\frac{1}{N^2} \sum_{i=1}^N \sum_{j=1}^N \left| \frac{1}{T} \sum_{t=1}^T [e_{it}e_{jt} - E(e_{it}e_{jt})] \right|^2 \right] = O_p(T^{-2}).$$

Given the above result, we have

$$\frac{1}{N} \sum_{i=1}^N \left| m_i' \hat{\Lambda} \hat{G} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T [e_t e_{it} - E(e_t e_{it})] \right|^2 = O_p\left(\frac{1}{NT}\right) + O_p\left(\frac{1}{T^2}\right).$$

This result, together with (B.18), gives

$$\frac{1}{N} \sum_{i=1}^N \mathcal{R}_i^2 = O_p\left(\frac{1}{NT}\right) + O_p\left(\frac{1}{T^2}\right). \quad (\text{B.19})$$

Notice that

$$\begin{aligned} \frac{1}{NT} \sum_{t=1}^T f_t e_t' \hat{\Sigma}_{ee}^{-1} M &= \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\hat{\sigma}_i^2} f_t e_{it} m_i' \\ &= \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} f_t e_{it} m_i' - \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{\hat{\sigma}_i^2 - \sigma_i^2}{\hat{\sigma}_i^2 \sigma_i^2} f_t e_{it} m_i'. \end{aligned}$$

The second term can be written as

$$\frac{1}{NT^2} \sum_{i=1}^N \sum_{t=1}^T \sum_{s=1}^T \frac{1}{\hat{\sigma}_i^2 \sigma_i^2} f_t e_{it} (e_{is}^2 - \sigma_i^2) m_i' + \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\hat{\sigma}_i^2 \sigma_i^2} \mathcal{R}_i f_t e_{it} m_i'$$

The second term of the above equation is bounded in norm by

$$C^5 \left[\frac{1}{N} \sum_{i=1}^N \|\mathcal{R}_i\|^2 \right]^{1/2} \left[\frac{1}{N} \sum_{i=1}^N \left\| \frac{1}{T} \sum_{t=1}^T f_t e_{it} \right\|^2 \right]^{1/2},$$

which is $O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2})$ by (B.19). The first term can be written as

$$\frac{1}{NT^2} \sum_{i=1}^N \sum_{t=1}^T \sum_{s=1}^T \frac{1}{\sigma_i^4} f_t e_{it} (e_{is}^2 - \sigma_i^2) m_i' - \frac{1}{NT^2} \sum_{i=1}^N \sum_{t=1}^T \sum_{s=1}^T \frac{\hat{\sigma}_i^2 - \sigma_i^2}{\hat{\sigma}_i^2 \sigma_i^4} f_t e_{it} (e_{is}^2 - \sigma_i^2) m_i'.$$

The first term of the above expression is $O_p(N^{-1/2}T^{-1})$. The second term is bounded in norm by

$$C^5 \left[\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2 \right]^{1/2} \left[\frac{1}{N} \sum_{i=1}^N \left\| \frac{1}{T} \sum_{t=1}^T f_t e_{it} \right\|^2 \cdot \left\| \frac{1}{T} \sum_{t=1}^T e_{it}^2 - \sigma_i^2 \right\|^2 \right]^{1/2},$$

which is $O_p(T^{-3/2})$. Given the above results, we have

$$\frac{1}{NT} \sum_{t=1}^T f_t e_t' \hat{\Sigma}_{ee}^{-1} M = \frac{1}{NT} \sum_{t=1}^T f_t e_t' \Sigma_{ee}^{-1} M + O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T^{3/2}}\right). \quad (\text{B.20})$$

Given (B.20), together with $\hat{R} = R + O_p(T^{-1/2})$, we immediately obtain (a). Given (B.20), together with $\hat{P} = P + O_p(T^{-1/2})$ and $\hat{\Lambda} = \Lambda + O_p(\frac{1}{\sqrt{NT}}) + O_p(\frac{1}{T})$, we also have (b).

We now consider (c). The left hand side of (c) is equal to

$$-\frac{1}{N} \sum_{i=1}^N \frac{\hat{\sigma}_i^2 - \sigma_i^2}{\hat{\sigma}_i^2 \sigma_i^2} m_i m_i' = -\frac{1}{N} \sum_{i=1}^N \frac{\hat{\sigma}_i^2 - \sigma_i^2}{\sigma_i^4} m_i m_i' + \frac{1}{N} \sum_{i=1}^N \frac{(\hat{\sigma}_i^2 - \sigma_i^2)^2}{\hat{\sigma}_i^2 \sigma_i^4} m_i m_i'.$$

We use i_1 and i_2 to denote the two expressions on the right hand side of the above equation. We first consider i_1 . Substituting (B.17) into this term, we obtain

$$\begin{aligned} i_1 &= -\frac{1}{N} \sum_{i=1}^N \frac{\hat{\sigma}_i^2 - \sigma_i^2}{\sigma_i^4} m_i m_i' = -\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^4} (e_{it}^2 - \sigma_i^2) m_i m_i' \\ &+ 2 \frac{1}{N} \sum_{i=1}^N \frac{1}{\sigma_i^4} \text{tr} \left[\hat{\Lambda} \hat{G} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T [e_t e_{it} - E(e_t e_{it})] m_i' \right] m_i m_i' - \frac{1}{N} \sum_{i=1}^N \frac{1}{\sigma_i^4} \mathcal{S}_i m_i m_i'. \end{aligned}$$

Consider the second expression. The (v, u) element of this expression $(v, u = 1, \dots, k)$ is

$$\text{tr} \left[\frac{1}{N} \sum_{i=1}^N \hat{\Lambda} \hat{G} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} \frac{1}{T} \sum_{t=1}^T [e_t e_{it} - E(e_t e_{it})] \frac{1}{\sigma_i^4} m_i' m_{iv} m_{iu} \right]$$

which can be proved to be $O_p(N^{-1}T^{-1/2}) + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2})$ similarly as Lemma B.3(a). The third term is bounded by

$$C^6 \left[\frac{1}{N} \sum_{i=1}^N \mathcal{S}_i^2 \right]^{1/2} = O_p(N^{-1}T^{-1/2}) + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2})$$

by (B.18). Hence, we have

$$i_1 = -\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^4} (e_{it}^2 - \sigma_i^2) m_i m_i' + O_p \left(\frac{1}{N\sqrt{T}} \right) + O_p \left(\frac{1}{\sqrt{NT}} \right) + O_p \left(\frac{1}{T^{3/2}} \right).$$

Proceed to consider i_2 . By

$$\hat{\sigma}_i^2 - \sigma_i^2 = \frac{1}{T} \sum_{t=1}^T (e_{it}^2 - \sigma_i^2) + \mathcal{R}_i,$$

we can write i_2 as

$$\frac{1}{N} \sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2 \sigma_i^4} \left[\frac{1}{T} \sum_{t=1}^T (e_{it}^2 - \sigma_i^2) \right]^2 m_i m_i' + 2 \frac{1}{N} \sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2 \sigma_i^4} \left[\frac{1}{T} \sum_{t=1}^T (e_{it}^2 - \sigma_i^2) \right] \mathcal{R}_i m_i m_i' + \frac{1}{N} \sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2 \sigma_i^4} \mathcal{R}_i^2 m_i m_i'.$$

We analyze the three terms at right-hand-side of the above equation one by one. The second term is bounded in norm by

$$2C^8 \left[\frac{1}{N} \sum_{i=1}^N \left| \frac{1}{T} \sum_{t=1}^T (e_{it}^2 - \sigma_i^2) \right|^2 \right]^{1/2} \left[\frac{1}{N} \sum_{i=1}^N \mathcal{R}_i^2 \right]^{1/2},$$

which is $O_p(N^{-1/2}T^{-1})$ by (B.19). The third term is bounded in norm by

$$C^8 \frac{1}{N} \sum_{i=1}^N \mathcal{R}_i^2 = O_p \left(\frac{1}{NT} \right) + O_p \left(\frac{1}{T^2} \right)$$

by (B.19). Finally, the first term can be written as

$$\frac{1}{N} \sum_{i=1}^N \frac{1}{\sigma_i^6} \left[\frac{1}{T} \sum_{t=1}^T (e_{it}^2 - \sigma_i^2) \right]^2 m_i m_i' - \frac{1}{N} \sum_{i=1}^N \frac{\hat{\sigma}_i^2 - \sigma_i^2}{\hat{\sigma}_i^2 \sigma_i^6} \left[\frac{1}{T} \sum_{t=1}^T (e_{it}^2 - \sigma_i^2) \right]^2 m_i m_i'$$

The first term of the above expression is equal to

$$\frac{1}{NT} \sum_{i=1}^N \frac{\kappa_{iA} - \sigma_i^4}{\sigma_i^6} m_i m_i' + O_p(N^{-1/2}T^{-1}).$$

The second term is bounded in norm by

$$C^{10} \left[\frac{1}{N} \sum_{i=1}^N (\hat{\sigma}_i^2 - \sigma_i^2)^2 \right]^{1/2} \left[\frac{1}{N} \sum_{i=1}^N \left| \frac{1}{T} \sum_{t=1}^T (e_{it}^2 - \sigma_i^2) \right|^4 \right]^{1/2} = O_p(T^{-3/2}).$$

Hence, we have

$$i_2 = \frac{1}{NT} \sum_{i=1}^N \frac{\kappa_{iA} - \sigma_i^4}{\sigma_i^6} m_i m_i' + O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T^{3/2}}\right).$$

Summarizing the results on i_1 and i_2 , we have (c). \square

PROOF OF THEOREM 4.2. We first derive the asymptotic behavior of A . Consider equation (A.16), using Lemma B.3 (a) and (f), Lemma B.5 (b) and Lemma B.4, we have

$$A + A' = \eta + \eta' + O_p(N^{-1}T^{-1/2}) + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2}),$$

where

$$\eta = \frac{1}{NT} \sum_{t=1}^T f_t e_t' \Sigma_{ee}^{-1} M \Lambda P^{-1}.$$

Let $\text{vech}(B)$ be the operation which stacks the elements on and below the diagonal of matrix B into a vector, for any square matrix B . Taking vech operation on both sides, we get

$$\text{vech}(A + A') = \text{vech}(\eta + \eta') + O_p(N^{-1}T^{-1/2}) + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2}).$$

Let D_r be the r -dimensional duplication matrix and D_r^+ be its Moore-Penrose inverse. By the basic fact that $\text{vech}(B + B') = 2D_r^+ \text{vec}(B)$, for any $r \times r$ matrix B , we have

$$2D_r^+ \text{vec}(A) = 2D_r^+ \text{vec}(\eta) + O_p(N^{-1}T^{-1/2}) + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2}). \quad (\text{B.21})$$

Furthermore, define

$$\zeta = \Lambda' \left[\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{m_i m_i'}{\sigma_i^4} (e_{it}^2 - \sigma_i^2) \right] \Lambda, \quad \mu = \Lambda' \left[\frac{1}{NT} \sum_{i=1}^N \frac{\kappa_{iA} - \sigma_i^4}{\sigma_i^6} m_i m_i' \right] \Lambda.$$

Proceed to consider equation (B.4). By Lemma B.5(c) and $\hat{\Lambda} - \Lambda = O_p(N^{-1/2}T^{-1/2}) + O_p(T^{-1})$ by Theorem 4.1, we have

$$\text{Ndg} \left\{ \hat{\Lambda}' \left(\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M \right) (\hat{\Lambda} - \Lambda) + (\hat{\Lambda} - \Lambda)' \left(\frac{1}{N} M' \hat{\Sigma}_{ee}^{-1} M \right) \hat{\Lambda} \right\}$$

$$= \text{Ndg}\{\zeta - \mu\} + O_p(N^{-1}T^{-1/2}) + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2}).$$

Using the same arguments in the derivation of (B.16), we have

$$\text{Ndg}(AP + PA') = \text{Ndg}(\zeta - \mu) + O_p(N^{-1}T^{-1/2}) + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2}).$$

Let $\text{veck}(B)$ be the operation which stacks the elements below the diagonal of matrix B into a vector, for any square matrix B . Let \mathcal{D} be the matrix such that $\text{veck}(B) = \mathcal{D}\text{vec}(B)$ for any $r \times r$ matrix B . By the preceding equation,

$$\text{veck}(AP + PA') = \text{veck}(\zeta - \mu) + O_p(N^{-1}T^{-1/2}) + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2}),$$

or equivalently

$$\mathcal{D}\text{vec}(AP + PA') = \mathcal{D}\text{vec}(\zeta - \mu) + O_p(N^{-1}T^{-1/2}) + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2}).$$

Using $\text{vec}(ABC) = (C' \otimes A)\text{vec}(B)$, we can rewrite the preceding equation as

$$\mathcal{D}[(P \otimes I_r) + (I_r \otimes P)K_r]\text{vec}(A) = \mathcal{D}\text{vec}(\zeta - \mu) + O_p(N^{-1}T^{-1/2}) + O_p(N^{-1/2}T^{-1}) + O_p(T^{-3/2}), \quad (\text{B.22})$$

where K_r is the r -dimensional communication matrix such that $K_r\text{vec}(B') = \text{vec}(B)$ for any $r \times r$ matrix B . By (B.21) and (B.22), we have

$$\begin{aligned} \left[\mathcal{D}[(P \otimes I_r) + (I_r \otimes P)K_r] \right] \text{vec}(A) &= \begin{bmatrix} 2D_r^+ \text{vec}(\eta) \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ \mathcal{D}\text{vec}(\zeta) \end{bmatrix} - \begin{bmatrix} 0 \\ \mathcal{D}\text{vec}(\mu) \end{bmatrix} \quad (\text{B.23}) \\ &+ O_p\left(\frac{1}{N\sqrt{T}}\right) + O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T^{3/2}}\right). \end{aligned}$$

Define

$$\mathbb{D}_1 = \begin{bmatrix} 2D_r^+ \\ \mathcal{D}[(P \otimes I_r) + (I_r \otimes P)K_r] \end{bmatrix}, \quad \mathbb{D}_2 = \begin{bmatrix} 2D_r^+ \\ 0_{\frac{1}{2}r(r-1) \times r^2} \end{bmatrix}, \quad \mathbb{D}_3 = \begin{bmatrix} 0_{\frac{1}{2}r(r+1) \times r^2} \\ \mathcal{D} \end{bmatrix}.$$

The above result can be rewritten as

$$\mathbb{D}_1\text{vec}(A) = \mathbb{D}_2\text{vec}(\eta) + \mathbb{D}_3\text{vec}(\zeta) - \mathbb{D}_3\text{vec}(\mu) + O_p\left(\frac{1}{N\sqrt{T}}\right) + O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T^{3/2}}\right). \quad (\text{B.24})$$

Also, notice that

$$\text{vec}(\eta) = \text{vec}\left[\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} f_t e_{it} m_i' \Lambda P^{-1}\right] = (P^{-1} \Lambda' \otimes I_r) \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} (m_i \otimes f_t) e_{it},$$

$$\text{vec}(\zeta) = \text{vec}\left[\Lambda' \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{m_i m_i'}{\sigma_i^4} (e_{it}^2 - \sigma_i^2) \Lambda\right] = (\Lambda \otimes \Lambda)' \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^4} (m_i \otimes m_i) (e_{it}^2 - \sigma_i^2)$$

and

$$\text{vec}(\mu) = \text{vec}\left[\Lambda' \frac{1}{NT} \sum_{i=1}^N \frac{\kappa_{iA} - \sigma_i^4}{\sigma_i^6} m_i m_i' \Lambda\right] = (\Lambda \otimes \Lambda)' \frac{1}{NT} \sum_{i=1}^N \frac{1}{\sigma_i^6} (m_i \otimes m_i) (\kappa_{iA} - \sigma_i^4).$$

Given the above three results, we can rewrite (B.24) as

$$\begin{aligned}
\text{vec}(A) &= \mathbb{D}_1^{-1} \mathbb{D}_2 (P^{-1} \Lambda' \otimes I_r) \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} (m_i \otimes f_t) e_{it} \\
&+ \mathbb{D}_1^{-1} \mathbb{D}_3 (\Lambda \otimes \Lambda)' \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^4} (m_i \otimes m_i) (e_{it}^2 - \sigma_i^2) \\
&- \mathbb{D}_1^{-1} \mathbb{D}_3 (\Lambda \otimes \Lambda)' \frac{1}{NT} \sum_{i=1}^N \frac{1}{\sigma_i^6} (m_i \otimes m_i) (\kappa_{iA} - \sigma_i^4) \\
&+ O_p\left(\frac{1}{N\sqrt{T}}\right) + O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T^{3/2}}\right).
\end{aligned} \tag{B.25}$$

Consider equation (A.18). Using the results of Lemma B.5 (a) and (b) and Lemma B.3 (e) and (f), we have

$$\begin{aligned}
\hat{\Lambda}' - \Lambda' &= -A' \Lambda' + \frac{1}{NT} \sum_{t=1}^T f_t e_t' \Sigma_{ee}^{-1} M R^{-1} + P^{-1} \Lambda' \frac{1}{NT} M' \Sigma_{ee}^{-1} \sum_{t=1}^T e_t f_t' \Lambda' \\
&+ O_p\left(\frac{1}{N\sqrt{T}}\right) + O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T^{3/2}}\right).
\end{aligned} \tag{B.26}$$

Notice that

$$\begin{aligned}
\text{vec}\left[\frac{1}{NT} \sum_{t=1}^T f_t e_t' \Sigma_{ee}^{-1} M R^{-1}\right] &= \text{vec}\left[\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} f_t e_{it} m_i' R^{-1}\right] \\
&= (R^{-1} \otimes I_r) \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} (m_i \otimes f_t) e_{it}
\end{aligned}$$

and

$$\begin{aligned}
\text{vec}\left[P^{-1} \Lambda' \frac{1}{NT} M' \Sigma_{ee}^{-1} \sum_{t=1}^T e_t f_t' \Lambda'\right] &= \text{vec}\left[P^{-1} \Lambda' \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} m_i e_{it} f_t' \Lambda'\right] \\
&= K_{kr} \text{vec}\left[\Lambda \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} f_t e_{it} m_i' \Lambda P^{-1}\right] \\
&= K_{kr} [(P^{-1} \Lambda') \otimes \Lambda] \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} (m_i \otimes f_t) e_{it},
\end{aligned}$$

where K_{mn} is the commutation matrix such that $K_{mn} \text{vec}(B) = \text{vec}(B')$ for any $m \times n$ matrix B .

Taking vectorization operation on the both sides of (B.26), we have

$$\begin{aligned}
\text{vec}(\hat{\Lambda}' - \Lambda') &= \left[K_{kr} [(P^{-1} \Lambda') \otimes \Lambda] + R^{-1} \otimes I_r \right] \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} (m_i \otimes f_t) e_{it} \\
&- K_{kr} (I_r \otimes \Lambda) \text{vec}(A) + O_p\left(\frac{1}{N\sqrt{T}}\right) + O_p\left(\frac{1}{\sqrt{NT}}\right) + O_p\left(\frac{1}{T^{3/2}}\right).
\end{aligned} \tag{B.27}$$

Substituting (B.25) into (B.27),

$$\begin{aligned} \text{vec}(\hat{\Lambda}' - \Lambda') &= \mathbb{B}_1 \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^2} (m_i \otimes f_t) e_{it} - \mathbb{B}_2 \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^4} (m_i \otimes m_i) (e_{it}^2 - \sigma_i^2) \\ &\quad + \frac{1}{T} \Delta + O_p \left(\frac{1}{N\sqrt{T}} \right) + O_p \left(\frac{1}{\sqrt{NT}} \right) + O_p \left(\frac{1}{T^{3/2}} \right), \end{aligned} \quad (\text{B.28})$$

where

$$\begin{aligned} \mathbb{B}_1 &= K_{kr}[(P^{-1}\Lambda') \otimes \Lambda] + R^{-1} \otimes I_r - K_{kr}(I_r \otimes \Lambda) \mathbb{D}_1^{-1} \mathbb{D}_2[(P^{-1}\Lambda') \otimes I_r], \\ \mathbb{B}_2 &= K_{kr}(I_r \otimes \Lambda) \mathbb{D}_1^{-1} \mathbb{D}_3(\Lambda \otimes \Lambda)', \\ \Delta &= \mathbb{B}_2 \frac{1}{N} \sum_{i=1}^N \sum_{t=1}^T \frac{1}{\sigma_i^6} (m_i \otimes m_i) (\kappa_{iA} - \sigma_i^4). \end{aligned}$$

Given the above results and by a Central Limit Theorem, we obtain as $N, T \rightarrow \infty$ and $N/T^2 \rightarrow 0$,

$$\text{sqr}tNT \left[\text{vec}(\hat{\Lambda}' - \Lambda') - \frac{1}{T} \Delta \right] \xrightarrow{d} N(0, \Omega),$$

where $\Omega = \lim_{N \rightarrow \infty} \Omega_N$ with

$$\Omega_N = \mathbb{B}_1(R \otimes I_r) \mathbb{B}_1' + \mathbb{B}_2 \left[\frac{1}{N} \sum_{i=1}^N \frac{\kappa_{iA} - \sigma_i^4}{\sigma_i^8} (m_i m_i') \otimes (m_i m_i') \right] \mathbb{B}_2'.$$

This completes the proof of Theorem 4.2. \square

PROOF OF THEOREM 4.5. By the definition of $\hat{f}_t = (\hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} M \hat{\Lambda})^{-1} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} z_t$ and A , we have

$$\hat{f}_t - f_t = -A' f_t + \hat{P}^{-1} \frac{1}{N} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} e_t.$$

From Corollary B.1, we know $A = O_p(\frac{1}{\sqrt{NT}}) + O_p(\frac{1}{T})$, then the first term of the above equation is $O_p(\frac{1}{\sqrt{NT}}) + O_p(\frac{1}{T})$. From Corollary A.1 (a)(b), we know $\hat{P} = P + o_p(1)$ and $\hat{P} = O_p(1)$, and from Assumption C.3, we know $P_\infty = \lim_{N \rightarrow \infty} P$ where P_∞ is positive definite matrix. Consider the part $\frac{1}{N} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} e_t$, which can be rewritten as

$$\frac{1}{N} \sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2} \hat{\Lambda}' m_i e_{it} = \frac{1}{N} \Lambda' M' \Sigma_{ee}^{-1} e_t - \frac{1}{N} \sum_{i=1}^N \frac{\hat{\sigma}_i^2 - \sigma_i^2}{\hat{\sigma}_i^2 \sigma_i^2} \Lambda' m_i e_{it} + \frac{1}{N} \sum_{i=1}^N \frac{1}{\hat{\sigma}_i^2} (\hat{\Lambda} - \Lambda)' m_i e_{it},$$

where m_i is the transpose of the i th row of M . Use a_1, a_2, a_3 to denote the three terms on the right hand side of the above equation. Term a_2 can be shown to be $O_p(\frac{1}{\sqrt{NT}}) + O_p(\frac{1}{T^{3/2}})$ by the equation (B.10). Term a_3 can be shown to be $O_p(\frac{1}{\sqrt{NT}}) + O_p(\frac{1}{T})$ by equation (A.18). Then we have

$$\frac{1}{N} \hat{\Lambda}' M' \hat{\Sigma}_{ee}^{-1} e_t = \frac{1}{N} \Lambda' M' \Sigma_{ee}^{-1} e_t + O_p \left(\frac{1}{\sqrt{NT}} \right) + O_p \left(\frac{1}{T} \right).$$

Therefore,

$$\hat{f}_t - f_t = P^{-1} \frac{1}{N} \Lambda' M' \Sigma_{ee}^{-1} e_t + O_p \left(\frac{1}{\sqrt{NT}} \right) + O_p \left(\frac{1}{T} \right)$$

Based on the above result, by a Central Limit Theorem, we obtain as $N, T \rightarrow \infty$ and $N/T^2 \rightarrow 0$,

$$\sqrt{N}(\hat{f}_t - f_t) \xrightarrow{d} N(0, P_{\infty}^{-1}).$$

This completes the proof of Theorem 4.5. \square

References

- Ahn, S.C. and Horenstein, A.R. (2013). Eigenvalue ratio test for the number of factors. *Econometrica*, 81 (3), 1203-1227.
- Anderson, T.W. (2003). An introduction to multivariate statistical analysis (3rd ed.). Hoboken, NJ: Wiley-Interscience.
- Bai, J. (2003). Inferential theory for factor models of large dimensions. *Econometrica*, 71 (1), 135-171.
- Bai, J. and Li, K. (2012). Statistical analysis of factor models of high dimension, *The Annals of Statistics*, 40 (1), 436-465.
- Bai, J. and Li, K. (2016). Maximum likelihood estimation and inference of approximate factor models of high dimension, *Review of Economics and Statistics*, 98(2), 298-309.
- Bai, J. and Ng, S. (2002). Determining the number of factors in approximate factor models. *Econometrica*, 70(1), 191-221.
- Bartlett, M.S. (1950). Test of significance in factor analysis. *British Journal of Psychology (Statistics Section)*, 3, 77-85.
- Carhart, M. M. (1997) On persistence of mutual fund performance. *Journal of Finance*, 52, 57-82.
- Chamberlain, G. and Rothschild, M. (1983). Arbitrage, factor structure, and mean-variance analysis on large asset markets. *Econometrica*, vol. 51(5), 1281-1304.
- Connor, G. and Korajczyk, R.A. (1986). Performance measurement with the arbitrage pricing theory : A new framework for analysis. *Journal of Financial Economics*, vol. 15(3), 373-394.
- Connor, G. and Korajczyk, R.A. (1988). Risk and return in an equilibrium APT: Application of a new test methodology. *Journal of Financial Economics*, vol. 21, 255-289.
- Connor, G. and O. Linton (2007). Semiparametric estimation of a characteristic-based factor model of stock returns. *Journal of Empirical Finance*, vol. 14, 694-717.
- Connor, G., M. Hagmann, and O. Linton (2012). Efficient semiparametric estimation of the Fama-French Model and extensions. *Econometrica*, vol. 80(2), 713-754.

- Doz, C., Giannone D., and Reichlin L. (2012). A quasi-maximum likelihood approach for large approximate dynamic factor models, *The Review of Economics and Statistics*, MIT Press, vol. 94(4), 1014-1024.
- Fama, E.F. and French, K.R. (1993). Common risk factors in the returns to stocks and bonds. *Journal of Financial Economics*, vol. 33, 3-56.
- Fan, J., Liao, Y., and Mincheva, M. (2011). High dimensional covariance matrix estimation in approximate factor models. *Annals of statistics*, 39(6), 3320.
- Fan, J., Liao, Y., and Mincheva, M. (2013). Large covariance estimation by thresholding principal orthogonal complements. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 75(4), 603-680.
- Fan, J., Liao Y. and Shi X. (2014). Risks of large portfolios, *Journal of Econometrics*, forthcoming.
- Fan, J., Liao Y. and Wang W. (2014). Projected principal component analysis in factor models, *manuscript*.
- Goyal, A. and Santa-Clara, P. (2003). Idiosyncratic risk matters, *Journal of Finance*, 58, 975-1008.
- Geweke, J. (1977). The dynamic factor analysis of economic time series, in: D.J. Aigner and A.S. Goldberger, eds., *Latent Variables in Socio-Economic Models*, (North-Holland, Amsterdam).
- Heckman, J.J., Stixrud, J., and Urzua, S. (2006). The effects of cognitive and noncognitive abilities on labor market outcomes and social behavior. *Journal of Labor Economics*, 24(3), 411-482.
- Jennrich, R.I. (1969). Asymptotic properties of non-linear Least Squares estimation. *Annals of Mathematical Statistics*, 40, 633-643.
- Kose, A., C. Otrok, and Whiteman, C. (2003). International business cycles: world region and country specific factors. *American Economic Review*, 93:4, 1216-1239.
- Lawley, D. N., and A. E., Maxwell (1971). *Factor analysis as a statistical method*. New York: American Elsevier.
- Matteson, D.S., M.W. McLean, D.B. Woodard and S.G. Henderson (2011). Forecasting emergency medical service call arrival rates. *Annals of Applied Statistics*, 5, 1379-1406.
- Onatski, A. (2010). Determining the number of factors from empirical distribution of eigenvalues, *Review of Economics and Statistics*, 92(4), 1004-1016.
- Rosenberg, B. (1974). Extra-market components of covariance in security prices, *Journal of Financial and Quantitative Analysis*, 9, 263-274.
- Ross, S. A. (1976). The arbitrage theory of capital asset pricing. *Journal of Economic Theory*, 13, 341-360.

- Sargent, T.J., and C.A. Sims (1977). Business cycle modeling without pretending to have too much a-priori economic theory, in: C. Sims et al., eds., *New Methods in Business Cycle Research* (Federal Reserve Bank of Minneapolis, Minneapolis).
- Stock, J.H. and Watson, M.W (2002). Forecasting using principal components from a large number of predictors, *Journal of the American Statistical Association*, 97, 1167-1179.
- Tsai, H. and Tsay, R. (2010). Constrained factor models, *Journal of the American Statistical Association*, vol. 105(492), 1593-1605.
- Zhou, Z. and Matteson, D.S. (2015). Temporal and spatio-temporal models for ambulance demand, book chapter in H.Yang and E.K.Lee, eds, *Healthcare Data Analytics*, Wiley Series in Operations Research and Management Science.